

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/042406

International filing date: 16 December 2004 (16.12.2004)

Document type: Certified copy of priority document


Document details: Country/Office: US  
Number: 60/531,809  
Filing date: 22 December 2003 (22.12.2003)

Date of receipt at the International Bureau: 03 February 2005 (03.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*January 21, 2005*

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/531,809

FILING DATE: *December 22, 2003*

RELATED PCT APPLICATION NUMBER: PCT/US04/42406



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

December 19, 1995

PTO/SB/16(11-95)

Approved for use through 01/31/98. OMB 0651-0037

Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

## PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT made under 37 CFR 1.53 (b)(2).

Docket Number

689290-180

Type a plus sign (+)  
inside this box →

+

## INVENTORS

LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Weigle Ebner	Bernd Reinhard		Hanau, Germany Gaithersburg, MD

## TITLE

PROSTATE-SPECIFIC PROTEINS EXPRESSED IN CANCER AND METHODS OF USE THEREOF

## CORRESPONDENCE ADDRESS

OLSTEIN, Elliot M. (Reg. No. 24,025); GILFILLAN, John G. (Reg. No. 22,746); BAIN, John N. (Reg. No. 18,651);  
LILLIE, Raymond J. (Reg. No. 31,778); SQUIRE, William (Reg. No. 25,378; GRANT, Alan J. (Reg. NO. 33,389);

Alan J. Grant, Esq., CARELLA, BYRNE, BAIN, GILFILLAN, CECCHI, STEWART & OLSTEIN, 6 Becker Farm Road, Roseland, NJ 07068.

STATE	ZIP CODE	COUNTRY
New Jersey	07068	USA

## ENCLOSED APPLICATION PARTS (check all that apply)



Specification

Number of Pages

56



Small Entity Statement



Drawing(s)

Number of Sheets



Other (specify)

## METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)



A check or money order is enclosed to cover the filing fees



The Commissioner is hereby authorized to charge  
filing fees and credit Deposit Account Number:

03-0678

FILING FEE

AMOUNT FOR SMALL ENTITY

\$80.00

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.



No.



Yes. The name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE

Alan J. Grant

DATE

22 December 2003

TYPE or PRINTED NAME

Alan J. Grant, Esq.

REGISTRATION NO.

33,389

☐ Additional inventors are being named on separately numbered sheets attached hereto

## USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

Burden Hour Statement: This form is estimated to take 2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, D.C. 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, DC 20231.

#52293 v1 - Prov.App.Cvr.Sht

EXPRESS MAIL NO: EL998937955US

22154 U.S. PTO  
60/531809

122203

# PROSTATE SPECIFIC PROTEINS EXPRESSED IN CANCER AND METHODS OF USE THEREOF

5

## FIELD OF THE INVENTION

10

The present invention relates to novel prostate specific transcripts expressed in most, if not all, prostate cancers, and potential uses in the screening, diagnosis, and treatment of prostate cancer, as well as the use of  
15 such transcripts for screening potential anti-cancer agents, including small organic compounds and other molecules, and development of therapeutic agents.

20

## BACKGROUND OF THE INVENTION

Cancer-linked genes are valuable in that they indicate genetic differences between cancer cells and normal cells, such as where a gene is expressed in a cancer cell but not in a non-cancer cell, or where said gene is over-expressed or  
25 expressed at a higher level in a cancer as opposed to normal or non-cancer cell. In addition, the expression of such a gene in a normal cell but not in a cancer cell, especially of the same type of tissue, can indicate important functions in the cancerous process. For example, screening assays for novel drugs are based on the response of model cell based systems *in vitro* to treatment with specific  
30 compounds. Such genes are also useful in the diagnosis of cancer and the identification of a cell as cancerous. Gene activity is readily measured by measuring the rate of production of gene products, such as RNAs and polypeptides encoded by such genes. Where genes encode enzyme proteins,

changes in the enzyme activity, including increased expression or change in activity due to mutation, are an indication of neoplastic activity. In addition, where said polypeptides represent cell surface antigens, available for detection or modulation through specific agents, such as antibodies, such as antibodies  
5 complexed with a cytotoxic agent, such as an apoptotic agent, the presence or absence of such antigen, or increased expression thereof, is useful for diagnosis and treatment of cancerous conditions. In accordance with the present invention, a cancer-linked gene has been identified and its putative amino acid sequence worked out. Such gene is over-expressed in prostate cancer cells and is useful in  
10 the diagnosing of cancer, the screening of anticancer agents and the treatment of cancer using such agents, especially in that this gene encodes an enzyme that provides a ready target for anti-tumor agents, including both antibodies that may bind to the protein as well as agents, such as small organic compounds, that modulate the activity of the enzyme, such as where the activity is correlated with  
15 cancerous characteristics and the agent serves to inhibit enzyme activity and thereby ameliorate, or prevent, the cancerous condition.

## **BRIEF SUMMARY OF THE INVENTION**

In accordance with the present invention, there is provided herein a cancer specific gene, linked especially to prostate cancer, or otherwise involved  
25 in the cancer initiating and facilitating process and the derived amino acid sequence thereof, including a polypeptide encoded by said gene (and the cDNA encoding said polypeptide), which polypeptide has cytochrome P450 oxidase activity.

30 In one aspect, the present invention relates to a process for identifying an agent that modulates the activity of a cancer-related gene comprising:

(a) contacting a compound with a cell containing a gene that corresponds to a polynucleotide having the sequence of SEQ ID NO: 2 or 4 and under conditions promoting the expression of said gene; and

(b) detecting a difference in expression of said gene relative to when said  
5 compound is not present  
thereby identifying an agent that modulates the activity of a cancer-related gene.

10 In various embodiments of such a process, the cell is a cancer cell and the difference in expression is a decrease in expression. Such polynucleotide may also include the sequence of SEQ ID NO: 2 or 4 (a cDNA) where a recombinant cell is used for screening, which cell has been genetically engineered to comprise and express said cDNA.

15 In another aspect, the present invention relates to a process for identifying an anti-neoplastic agent comprising contacting a cell exhibiting neoplastic activity with a compound first identified as a cancer related gene modulator using an assay process disclosed herein and detecting a decrease in said neoplastic activity after said contacting compared to when said contacting does not occur.

20

The present invention also relates to a process for identifying an anti-neoplastic agent comprising administering to an animal exhibiting a cancer condition an effective amount of an agent first identified according to a process of one of one of the assays disclosed according to the invention and detecting a  
25 decrease in said cancerous condition.

The present invention further relates to a process for determining the cancerous status of a cell, comprising determining an increase in the level of expression in said cell of at least one gene that corresponds to a polynucleotide  
30 having the sequence of SEQ ID NO: 2 or 4 and wherein an elevated expression

relative to a known non-cancerous cell indicates a cancerous state or potentially cancerous state.

5       The present invention additionally relates to an isolated polypeptide,  
10    encoded by one of the polynucleotide transcripts disclosed herein, comprising an  
      amino acid sequence homologous to the amino acid sequence of SEQ ID NO: 3  
      or 5 wherein any difference between said amino acid sequence and the  
      sequence of SEQ ID NO: 3 or 5 is due solely to conservative amino acid  
      substitutions and wherein said isolated polypeptide comprises at least one  
      immunogenic fragment.

      The present invention also relates to an antibody that reacts with a  
      polypeptide as disclosed herein, preferably a polypeptide comprising the amino  
      acid sequence of SEQ ID NO: 3 or 5. Such an antibody may be polyclonal,  
15    monoclonal, recombinant or synthetic in origin.

      In one such embodiment, said antibody is associated, either covalently  
      or non-covalently, with a cytotoxic agent, for example, an apoptotic agent.  
      Thus, the present invention relates to an immunoconjugate comprising an  
20    antibody of the invention and a cytotoxic agent.

      The present invention also relates to a process for treating cancer  
      comprising contacting a cancerous cell with an agent having activity against an  
      expression product encoded by a gene sequence corresponding to that of SEQ  
25    ID NO: 2 or 4. In one such embodiment, the cancerous cell is contacted *in vivo*.

      The present invention further encompasses an immunogenic composition  
      comprising a polypeptide disclosed herein, as well as compositions formed using  
      antibodies specific for these polypeptides.

30

The present invention is also directed to uses of such compositions. Such uses include a method for treating cancer in an animal afflicted therewith comprising administering to said animal an amount of an immunogenic composition of one or more of the polypeptides disclosed herein where such amount is an amount sufficient to elicit the production of cytotoxic T lymphocytes specific for a polypeptide of the invention, preferably a polypeptide incorporating the sequence of SEQ ID NO: 3 or 5. In a preferred embodiment, the animal to be so treated is a human patient.

## DEFINITIONS

As used herein, the terms "portion," "segment," and "fragment," when used in relation to polypeptides, refer to a continuous sequence of residues, such as amino acid residues, which sequence forms a subset of a larger sequence. For example, if a polypeptide were subjected to treatment with any of the common endopeptidases, such as trypsin or chymotrypsin, the oligopeptides resulting from such treatment would represent portions, segments or fragments of the starting polypeptide. When used in relation to a polynucleotides, such terms refer to the products produced by treatment of said polynucleotides with any of the common endonucleases.

As used herein, the term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). It could also be produced recombinantly and subsequently purified. For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides, for example, those prepared recombinantly, could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated



in that such vector or composition is not part of its natural environment. In one embodiment of the present invention, such isolated, or purified, polypeptide is useful in generating antibodies for practicing the invention, or where said antibody is attached to a cytotoxic or cytolytic agent, such as an apoptotic agent.

5

The term "percent identity" or "percent identical," when referring to a sequence, means that a sequence is compared to a claimed or described sequence after alignment of the sequence to be compared (the "Compared Sequence") with the described or claimed sequence (the "Reference Sequence").

10 The Percent Identity is then determined according to the following formula:

$$\text{Percent Identity} = 100 [1 - (C/R)]$$

wherein C is the number of differences between the Reference Sequence and the  
15 Compared Sequence over the length of alignment between the Reference Sequence and the Compared Sequence wherein (i) each base or amino acid in the Reference Sequence that does not have a corresponding aligned base or amino acid in the Compared Sequence and (ii) each gap in the Reference Sequence and (iii) each aligned base or amino acid in the Reference Sequence that is different  
20 from an aligned base or amino acid in the Compared Sequence, constitutes a difference; and R is the number of bases or amino acids in the Reference Sequence over the length of the alignment with the Compared Sequence with any gap created in the Reference Sequence also being counted as a base or amino acid.

25

If an alignment exists between the Compared Sequence and the Reference Sequence for which the percent identity as calculated above is about equal to or greater than a specified minimum Percent Identity then the Compared Sequence has the specified minimum percent identity to the Reference Sequence even  
30 though alignments may exist in which the hereinabove calculated Percent Identity is less than the specified Percent Identity.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide.

5

In accordance with the present invention, the term "DNA segment" or "DNA sequence" refers to a DNA polymer, in the form of a separate fragment or as a component of a larger DNA construct, which has been derived from DNA isolated at least once in substantially pure form, i.e., free of contaminating endogenous materials and in a quantity or concentration enabling identification, manipulation, and recovery of the segment and its component nucleotide sequences by standard biochemical methods, for example, using a cloning vector. Such segments are provided in the form of an open reading frame uninterrupted by internal nontranslated sequences, or introns, which are typically present in eukaryotic genes. Sequences of non-translated DNA may be present downstream from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

The term "coding region" refers to that portion of a gene which either naturally or normally codes for the expression product of that gene in its natural genomic environment, i.e., the region coding *in vivo* for the native expression product of the gene. The coding region can be from a normal, mutated or altered gene, or can even be from a DNA sequence, or gene, wholly synthesized in the laboratory using methods well known to those of skill in the art of DNA synthesis.

25

In accordance with the present invention, the term "nucleotide sequence" refers to a heteropolymer of deoxyribonucleotides. Generally, DNA segments encoding the proteins provided by this invention are assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being expressed in a recombinant

30

transcriptional unit comprising regulatory elements derived from a microbial, eukaryotic or viral operon.

5 The term "expression product" means that polypeptide or protein that is the natural translation product of the gene and any nucleic acid sequence coding equivalents resulting from genetic code degeneracy and thus coding for the same amino acid(s).

10 The term "active fragment," when referring to a coding sequence, means a portion comprising less than the complete coding region whose expression product retains essentially the same biological function or activity as the expression product of the complete coding region.

15 The term "primer" means a short nucleic acid sequence that is paired with one strand of DNA and provides a free 3'-OH end at which a DNA polymerase starts synthesis of a deoxyribonucleotide chain.

20 The term "promoter" means a region of DNA involved in binding of RNA polymerase to initiate transcription. The term "enhancer" refers to a region of DNA that, when present and active, has the effect of increasing expression of a different DNA sequence that is being expressed, thereby increasing the amount of expression product formed from said different DNA sequence.

25 The term "open reading frame (ORF)" means a series of triplets coding for amino acids without any termination codons and is a sequence (potentially) translatable into protein.

30 As used herein, reference to a "DNA sequence" includes both single stranded and double stranded DNA. Thus, the specific sequence, unless the context indicates otherwise, refers to the single strand DNA of such sequence,

the duplex of such sequence with its complement (double stranded DNA) and the complement of such sequence.

As used herein, "corresponding genes" refers to genes that encode an  
5 RNA that is at least 90% identical, preferably at least 95% identical, most preferably at least 98% identical, and especially identical, to an RNA encoded by one of the nucleotide sequences disclosed herein (i.e., SEQ ID NO: 1, 2, 3 and 4). Such genes will also encode the same polypeptide sequence as any of the sequences disclosed herein, preferably SEQ ID NO: 1, 2, 3 and 4, but may  
10 include differences in such amino acid sequences where such differences are limited to conservative amino acid substitutions, such as where the same overall three dimensional structure, and thus the same antigenic character, is maintained. Thus, amino acid sequences may be within the scope of the present invention where they react with the same antibodies that react with polypeptides  
15 comprising the sequences of SEQ ID NO: 5 and 6. A "corresponding gene" includes splice variants thereof.

The genes identified by the present disclosure are considered "cancer-related" genes, as this term is used herein, and include genes expressed at  
20 higher levels (due, for example, to elevated rates of expression, elevated extent of expression or increased copy number) in cancer cells relative to expression of these genes in normal (i.e., non-cancerous) cells where said cancerous state or status of test cells or tissues has been determined by methods known in the art, such as by reverse transcriptase polymerase chain reaction (RT-PCR) as  
25 described in the Examples herein. In specific embodiments, this relates to the genes whose sequences correspond to the sequences of SEQ ID NO: 2 and 4.

As used herein, the term "conservative amino acid substitutions" are defined herein as exchanges within one of the following five groups:  
30

1. Small aliphatic, nonpolar or slightly polar residues:

Ala, Ser, Thr, Pro, Gly;

II. Acids or amides:

Asp, Asn, Glu, Gln;

III. Basic residues:

5 His, Arg, Lys;

IV. Aliphatic residues:

Met Leu, Ile, Val, Cys

V. Aromatic residues:

10 Phe, Tyr, Trp

## DETAILED SUMMARY OF THE INVENTION

15 The present invention relates to processes for utilizing a nucleotide sequence for a cancer-linked gene, polypeptides encoded by such sequences and antibodies reactive with such polypeptides in methods of treating and diagnosing cancer, preferably prostate cancer, and in carrying out screening assays for agents effective in reducing the activity of cancer-linked genes and  
20 thereby treating a cancerous condition.

In accordance with the present invention, the sequences of the polynucleotides and polypeptides of the invention are as follows:

25 SEQ ID NO: 1 - AF109300

GCCTCATCCCTGGGTTGTGGTGTGGACATTGTGGGTGTCTCCACAGGAGCC  
CCAGGGCCACGAAAGCTGGGGTGGCCTCTGCCCTTCTGGGGTTCTTTTC  
CTGCACAGCTGCTTTCTGACTCCACCCACAGCTGGGAGCAGGTGCCGGAG  
30 CCCCggCCTGCATGGCCCTGTGAAGGCCACTCTGGGCGTTTGGGTGGGCG  
TGAGTGCCTTCCTCTGCTCCAGCATGTGGTTTTCTCCGTTGGCCGCCTCC

TGGACCTCCTGGTGCCTGACATCCCAGAGTCTGTGGAGATCAAAGTGAAGC  
 GGGAGTACTACCTGGCTAAGCAGGCACTGGCTGAGAATGAGGTTCTTTTTG  
 GAACGAACGGAACAAAGGATGAGCAGCCGAGGGCTCAGAGCTCAGCTCC  
 CACTGGACACCCCTTACGCTTCCCAAGGCCAGCCAGCTGCAGCAGTGACG  
 5 CCTGGAAGGACATCTGGTGGTCTTAGGGGAGTGGCCCTCCTGAGCCCT  
 GCGAGCAGCGTCTTTTCTCTTCCCTCAGGCAGCGGCTGTGTGAACCGCT  
 GGCTGCTGTTGTGCCTCATCTCTGGGCACATTGCTGCTTCCCCCAGCGC  
 CGGCTTCTCTCCTCAGAGCGCCTGTCACTCCATCCCCGGCAGGGAGGGAC  
 CGTCAGCTCACAAGGCCCTCTTTGTTTCTGCTCCCAGACATAAGCCCAAG  
 10 GGGCCCTGCACCCAAGGGACCCTGTCCCTCGGTGGCTCCCAGGCCCC  
 TGGACACGACAGTTCTCCTCAGGCAGGTGGGCTTTGTGGTCTCGCCGCC  
 CCTGGCCACATCGCCCTCTCCTCTTACACCTGGTGACCTTGAATGTTTCAG  
 AGCGCAGGGCGGTTCTCCCTCGTGTCTCTGGACCCACCCGCCCTTCTCT  
 GCCCTGTTTGCGCAGGGACATCACCCACATGCCCCAGCTCTCGGACCCTG  
 15 CAGCTCTGTGTCCAGGCCACAGCAAAGGTCTGTTGAACCCCTCCCTCCAT  
 TCCAGTTATCTGGGTCTCTGGATTCTTCTGTTTCTGAATCAGGCTCTGC  
 TTTCCCCCTAGCCACTACAGGCAGCCTCTGACAGTGCCGCTTTACTTGCATT  
 CTGCAGCAATTACATGTGTCTTTTGATCCTTGCCCAACTTCCCTCCCTCTC  
 CCAGCTCCTGGCCCCCTGGCCCAGGGCCCCCTCTTGCTGTTTTTACCTCTGT  
 20 CCTTGGGGCCTAGTACCCAGCAAGCACCCAAATGGGGGAGGTTTTGGGAT  
 GAGAGGAGGAAACGTGTATACCTGTAACATCTGGTGGCTCTTCCCCAGAA  
 GTTTGTGTTACATATAATTGTTTTCCACGCTGGATCATAATGTGACGTGCA  
 GTTCTGCCCTGTGCTGGGGAGCCACATGAAGCTTCCCTGGCTAACTTGCT  
 ACCCCGCAGCAATCCCAGTGTGGCTGTCTGCTTGCTAAAAAATGGATCTGT  
 25 GCTCATCTGTATTGATGTCTTGAGTTCTACAAGTGGAACCTAAGTGTCAA  
 AAAGAATATGTGGTTTTTAGGGGATCCTCTAGAGTCGACCT

SEQ ID NO: 2 – PCa3 RACE

30

caaggtgagggcatgcgaatggctgccaactgcctggcggggctccaagggccacccctccccacctctgtc  
 ccgcatgtagggacgggactctactgccagaccaggctcacgctgagaggtgggccaatgacctccgagacct  
 ctccggaagccactgtgccaggatgctgcggcgacgggcccaggaaagagacagacacgtctctgatcgat  
 gtgagccccctgaggcagagaagaggggctcttacgggagcacagcccacgcctcggagccagggtggac  
 5 agcaagcggccgcctgcagagctgggagtcctgccaagccccggatcgactctgctcgtttgggagagg  
 acctgaagctagacaggcagcaggacagtgccgccccgggacagaacagacatgcacaggacctggcggg  
 agactttctggataatctctgtgcggctgggctgtgtgtagaccagcaggacgtccaggacgggaacaccaca  
 gtgactacgcctctcctcgcgcctcctgggctgtgtctgtactacgcgaagacctgcgcctgaagctgcc  
 ttgaggaggttaccacaccaggcctccaactggtgcggcgctgctggatggctgggcatcccaacgtct  
 10 gctggaggtgtgcccagacgtacccccgagtactactcctgccggttcagagtgaacaagctgccacgttct  
 cgggagtgacaaccaggacaccttctcaagaccaagagccaccaaatctgttgagatcctggccaag  
 acccgtatggccacgagaagaaaaacctgctggatccaaccagctgtcggcagagg gttctctcag  
 tgccgccttc ccctgcatg acgccccct caagacgccc ccaggggcc cgcaggctcc  
 acgcctcaac cagcgccaag tcttttcca gactgggcg cgtggggca agtgaacaa  
 15 gtaccagccc ctgaccacg tgcgaggtat ctgggggag aaggtggccctactctgc ctgctcggg  
 ttttacag gctggctct gccagcggca gtgtgggca cactgtgtt cctgtgggc tgttctgg  
 tgttctcaga cataccacg caggaactgt gtggcagcaa ggacagcttc gagatgtgcc cacttgcct  
 cgactgcct ttctgctgc tctccagcg cgtgcctg gccaggccggccgctg ttgaccacg  
 gcggcaccgt gttctcagc ttgtcatg cactgtgggc cgtgctgct ctggagtact ggaagcggaa  
 20 gagcgccacg ctggcctacc gctgggactg ctctgactac gaggacactg aggagaggcc  
 toggccccag ttgccgcct cagccccat gacagccccg aacccatca cgggtgagga  
 cgagccctac ttccctgaga ggagccgcg gcgcgcgatg ctggccggct ctgtgtgat cgtgtgatg  
 gtggccgtg ttgtcatgt cctgtgtct atcatcctgt accgtgcat catggccatc gtgtgtcca  
 ggtggggcaa cacccttc gcagcctgg cctctgcac gccagcctc accgggtctg tagtgaacct  
 25 cgtcttcac ctatcctct ccaagatcta tgtatccctg gccacgtcc tgacacgatg gaaatgcac  
 cgcaccaga ccaagtcca ggacgcctc accccaagg tttcatctt ccagtctgc aacttctact  
 cctcaccgt ctacattgcc ttctcaagg gcaggttgt gggatacca ggaactacc acacctgtt  
 tggagtccg aatgaggagtgcgcggctg gaggtgcct gatcgagctg gcacaggagc tctgtgtat  
 catgtgggc aagcaggtca tcaacaacat gcaggaggtc ctatccc ga agctaaaggg  
 30 ctgtgtgcag aagttccgc ttgcctcaa gaagaggaag gcgggagctt ctgcaggggc  
 tagccagggg cctgggagg acgactatga gctgtgccc tgtgagggtc tgttgacga gtacctgaa

atggtgtcgcagt tgggttcgt caccatcttc gtggcgcct gtccgcctgc gccgccttc gccctgtca  
 acaactgggt ggagatcgc ttggacgcg gcaagttcgt ctgcgagtac cggcgccgg  
 tggcgagcg cggccaggac atcgcatct ggtccacat cctggcggc ctacgcacc tggcggtat  
 cagcaacgcc ttctcctg cctctcgtc cgactctg ccgcgcct actaccggt gaccgcgccc  
 5 cagcactgc gggcctct caactcacg ctggcgag cccgctct ctccgcgc gcgcacaacc  
 gcacgtgcag ttagcagga cgagtgcag acagaactct cagacacgg attaagaagg  
 aagaggtttt  
 ttattcgccggggcgctggcagactgtgtcttcagagcggagctgcggaataaattcttagcccttg  
 aagggcttacaactcaagggtctacgtgaagagtcataatagatcaa gtatcgggt ttccgggat  
 10 acgatggaca ttattccag acctactgga atctcttc catccgctg gccctgtca ttgtttga  
 ggtagccgag gcacctcgt gtctccat ccatggcatg agggccgc cctgtctt gcctaattg  
 agcagtg gtgagg ggtcgtgc gtcactct gctgtgtat ctgtgtaaa tcagagctct  
 tctcgcacc tgcgtttcc ctgcctggc tcactcctg gtgtgtgt ggacattgt ggtgtctca  
 caggagcccc agggccaaga aagctgggt ggcctctgc cctctggg ttctttcc tgcacagctg  
 15 ctctgact ccaccacag ctgggagcag gtgcggagc ccggcctgc ctggcctgt gaagccact  
 ctggcggtt ggtggggt gagtgctc ctgtctcc agcatgtgt ttctcgtt gccgcctcc  
 tggactctt ggtcctgac atccagagt ctgtggagt caaagtgaag cgggagctact acctggcta  
 gcaggcactg ctgagaatg ag gtct tttggaac aacggaaca aggatagca ccccgagggc  
 tcagag ct agtccact ggacacct cagggtcc aaggccagc agctgcagca gtgacgctg  
 20 gaaggacatc tgggtgtct taggggagtg gcccctct agccctcga gcagctct ttctcttc  
 cctcaggcag cggctgtgt aaccgctgc tctgtgtg cctatctt ggcacattg cctgtctcc  
 ccagcgcg cctctctc tcagagcgc tgcactca tcccggag ggaggagcgc tcagctca  
 aggcctct ttgtctgc tccagacat aagccaagg ggcctgca ccaaggac cctgtctc  
 ggtggcct ccaggcccc ggacacgaca gtctctca ggcaggtgg cttgtgtc ctgcgcgc  
 25 ctggccacat cgccctctc tctacact ggtgacct gaattgtta gagcgagg cgtctctc  
 tctgtctc tggaccacc cgcccttc tccctgtt gcgaggag ataccaca tccccagct  
 ctggacct cgactctgt gtccaggcc acagcaaagg tctgtgaac cctctctc attccagtt  
 atctgggtc tctgattct tctgttct gaatcaggt ctgtctcc ctagccact acaggcagc  
 tctgacagt ccgtttact tgcattct agcaattaca tgtctctt tgcctctg ccaactcc  
 30 tccctccc agtctctgc cctggccca gggccctct tctgtttt accttttc ctggggcct  
 agtaccagc aagcaccaca atgggggagg ttgggatg agaggaggaa acgtgtatac



ctgtaacatc tgggtgctct tccccagaa gttgtgttc atacataatt gtttccacg ctggatcata  
 atgtgacgtg cagttctgcc ctgtgctggg gagccacatg aagcttcccc tggctaactt gctaccccg  
 agcaatccca gtgtggcgt ctgctgtcta aaaaatggat ctgtgctcat ctgtaaaaaa aaaaaaaaaa  
 aaaaaaaa

5

SEQ ID NO: 3 – PCa3\_protein

QGEGRMAATAWAGLQGPPLPTLCPAVRTGLYCRDQA  
 HAERWAMTSETSSGSHCARMLRRRAQEEDSTVLIDVS  
 10 PPEAEKRGSYGSTAHASEPGGQQAACRAGSPAKPRI  
 DFLVWVEEDLKLDLRQQDSAARDRTDMHRTWRETFDND  
 LRAAGLCVDQDQDVQDGNNTTVHYALLSASWAVLCYYAE  
 DLRLKLPLQELPNQASNWSAGLLAWLGIPNVLLLEVVPD  
 VPPEYYSCRFRVNLKPRFLGSDNQDTFFTSTKRHQILF  
 15 EILAKTPYGHEKKNLLGIHQLLAEGVLSAAFPLHDGPFK  
 TPPEGPPAPRLNQRQVLFQHWARWGKWNKYQPLDHV  
 RRYFGEKVALYFAWLGFYTGWLLPAAVVGTLVFLVGCFL  
 LVFSDIPTQELCGSKDSFEMCPLCLDCPFWLLSSACAL  
 AQAGRLFDHGGTVFFSLFMALWAVLLLEYWKRKSATLA  
 20 YRWDCSDYEDTEERPRPQFAASAPMTAPNPITGEDEP  
 YFPERSRARRMLAGSVVIVVMVAVVVMCLVSIILYRAIM  
 AIVVSRSGNTLLAAWASRIASLTGSVNVNLVFILILSKIYV  
 SLAHVLTRWEMHRTQTKFEDAFTLKVFIFQFVNFYSSP  
 VYIAFFKGRFVGYPGNYHTLFGVRNEECAAGGCLIELA  
 25 QELLVIMVGKQVINNMQEVLPCEGLFDEYLEMVLQFG  
 KAGASAGASQGPWEDDYELVPCEGLFDEYLEMVLQFG  
 FVTIFVAACPLAPLALLNNWVEIRLDARKFVCEYRRPV  
 AERAQDIGIWFHILAGLTHLAVISNAFLAFSSDFLPRA  
 YYRWTRAHDLRGFLNFTLARAPSSFAAAHNRTCSVAG  
 30 RVADRTPTQTPD

SEQ ID NO: 4 – PCa3\_splice variant

	CAAG	GTGAGGGCAT	GCGAATGGCT	GCCACTGCCT	GGGCGGGGCT
5	CCAAGGGCCA	CCCCTCCCCA	CCCTCTGTCC	CGCAGTGAGG	
	ACGGGACTCT	ACTGCCGAGA	CCAGGCTCAC	GCTGAGAGGT	
	GGGCCATGAC	CTCCGAGACC	TCTTCGGAA	GCCACTGTGC	CAGG
	ATGCTGCGGC	GACGGGCCCCA	GGAAGAGGAC	AGCACCCTCC	
	TGATCGATGT	GAGCCCCCCT	GAGGCAGAGA	AGAGGGGGCTC	
10	TTACGGGAGC	ACAGCCACG	CCTCGGAG	CCAGGTGGAC	AGCAAGCGGC
	CGCCTGCAGA	GCTGGGAGTC	CTGCCAAGCC	CCGGATCGAC	
	TTCGTCCTCG	TTTGGGAGGA	GGACCTGAAG	CTAGACAGGC	
	AGCAGGACAG	TGCCGCCCCG	GACAGAACAG	ACATGCACAG	
	GACCTGGCGG	GAGACTTTTC	TGGATAATCT	TCGTGCGGCT	
15	GGGCTGTGTG	TAGACCAGCA	GGACGTCCAG	GACGGGAACA	
	CCACAGTGCA	CTACGCCCTC	CTCAGCGCCT	CCTGGGCTGT	
	GCTCTGCTAC	TACGCCGAAG	ACCTGCGCCT	GAAGCTGCCC	
	TTGCAGGAGT	TACCCAACCA	GGCCTCCAAC	TGGTCGGCCG	
	GCCTGCTGGC	ATGGCTGGGC	ATCCCCAACG	TCCTGCTGGA	
20	GGTTGTGCCA	GACGTACCCC	CCGAGTACTA	CTCCTGCCGG	
	TTCAGAGTGA	ACAAGCTGCC	ACGCTTCCTC	GGGAGTGACA	
	ACCAGGACAC	CTTCTTCACA	AGCACCAAGA	GGCACCAAAT	TCTGTTTGAG
	ATCCTGGCCA	AGACCCCGTA	TGGCCACGAG	AAGAAAAACC	
	TGCTTGGGAT	CCACCAGCTG	CTGGCAGAGG	GTGTCCTCAG	
25	TGCCGCCTTC	CCCCTGCATG	ACGGCCCCCTT	CAAGACGCCC	
	CCAGAGGGCC	CGCAGGCTCC	ACGCCTCAAC	CAGCGCCAAG	
	TCCTTTTCCA	GCACTGGGCG	CGCTGGGGCA	AGTGAACAA	
	GTACCAGCCC	CTGGACCACG	TGCGCAGGTA	CTTCGGGGAG	
	AAGGTGGCCC	TCTACTTCGC	CTGGCTCGGG	TTTACACAG	
30	GCTGGCTCCT	GCCAGCGGCA	GTGGTGGGCA	CACTGGTGTT	
	CCTGGTGGGC	TGCTTCCTGG	TGTTCTCAGA	CATACCCACG	

	CAGGAACTGT	GTGGCAGCAA	GGACAGCTTC	GAGATGTGCC
	CACTTTGCCT	CGACTGCCCT	TTCTGGCTGC	TCTCCAGCGC
	CGCCAGG	CCGGCCGGCTG	TTCGACCACG	GCGGCACCGT
	TTGTTTCATGG	CACTGTGGGC	CGTGCTGCTG	CTGGAGTACT
5	GGAAGCGGAA	GAGCGCCACG	CTGGCCTACC	GCTGGGACTG
	CTCTGACTAC	GAGGACACTG	AGGAGAGGCC	TCGGCCCCAG
	TTTGCCGCT	CAGCCCCCAT	GACAGCCCCG	AACCCCATCA
	CGGGTGAGGA	CGAGCCCTAC	TTCCCTGAGA	GGAGCCGCGC
	GCGCCGCATG	CTGGCCGGCT	CTGTGGTGAT	CGTGGTGATG
10	GTGGCCGTGG	TGGTCATGTG	CCTCGTGTCT	ATCATCCTGT
	ACCGTGCCAT	CATGGCCATC	GTGGTGCCA	GGTCGGGCAA
	CACCCCTTCTC	GCAGCCTGGG	CCTCTCGCAT	CGCCAGCCTC
	ACGGGGTCTG	TAGTGAACCT	CGTCTTCATC	CTCATCCTCT
	CCAAGATCTA	TGTATCCCTG	GCCACGTCC	TGACACGATG
	GGAAATGCAC	CGCACCAGA	CCAAGTTCGA	GGACGCCTTC
15	ACCCTCAAGG	TGTTTCATCTT	CCAGTTCGTC	AACCTTCTACT
	CCTCACCCGT	CTACATTGCC	TTCTTCAAGG	GCAGGTTTGT
	GGGATACCCA	GGCAACTACC	ACACCTTGTT	TGGAGTCCGC
	GATCGAGCTG	AATGAGGAGT	GCGCGGCTG	GAGGCTGCCT
	GATCGAGCTG	GCACAGGAGC	TCCTGGTCAT	CATGGTGGGC
20	AAGCAGGTCA	TCAACAACAT	GCAGGAGGTC	CTCATCCC
	GAAGCTAAAGGG	CTGGTGGCAG	AAGTTCGGC	TTCGCTCCAA
	GAAGAGGAAG	GCGGGAGCTT	CTGCAGGGC	TAGCCAGGGG
	CCCTGGGAGG	ACGACTATGA	GCTTGTGCCC	TGTGAGGGTC
	TGTTTGACGA	GTACCTGGAA	ATGGTGCTGCAGT	TCGGCTTCGT
	GTGGCCGCTC	GTCCGCTCGC	GCCGCTCTTC	GCCCTGCTCA
25	ACAACTGGGT	GGAGATCCGC	TTGGACGCGC	GCAAGTTCGT
	CTGCGAGTAC	CGGCGCCCGG	TGGCCGAGCG	CGCCAGGAC
	ATCGGCATCT	GGTTCCACAT	CCTGGCGGGC	CTCACGCACC
	TGGCGGTCTAT	CAGCAACGCC	TTCTCTCTGG	CCTTCTCGTC
	CGACTTCTGC	CCGCGCGCCT	ACTACCGGTG	GACCCGCGCC
30	CACGACCTGC	GCGGCTTCCT	CAACTTCACG	CTGGCGCGAG
	CCCCGTCCTC			

	CTTCGCCGCC	GCGCACAACC	GCACGTGCAG	GTATCGGGCT
	TTCCGGGATG	ACGATGGACA	TTATTCCCAG	ACCTACTGGA
	ATCTTCTTGC	CATCCGCCCTG	GCCTTCGTCA	TTGTGTTTGA
	GGTAGCCGAG	GCACCTGCTG	GTTCTCCCAT	CCATGGCATG
	AGGCCCCGAC	5	CCTGTGCTTT	GCCTAATTCG
	AGCACGTG	GTGAGG	GGTCGGTGCC	GTCACCTCCT
	GCTGTGTCAT	CTTGGTCAAA	TCAGAGCTCT	TCTCTGCACC
	TGCGTTTTCC	CTGCCTGGCC	TCATCCCTGG	GTTGTGGTGT
	GGACATTGTG	GGTGTCTCCA	CAGGAGCCCC	AGGGCCACGA
	AAGCTGGGGT	GGCCTCTGCC	CCTTCTGGGG	TTCTTTTCC
	TGCACAGCTG	CTTTCTGACT	10	CCACCCACAG
	CTGGGAGCAG	GTGCCGGAGC	CCCGGCCTGC	CTGGCCCTGT
	GAAGGCCACT	CTGGGCGTTT	GGGTGGGCGT	GAGTGCCTTC
	CTCTGCTCCC	AGCATGTGGT	TTTCTCCGTT	GGCCGCCTCC
	TGGACCTCCT	GGTGCCTGAC	ATCCCAGAGT	CTGTGGAGAT
	CAAAGTGAAG	CGGGAGTACT	ACCTGGCTAA	GCAGGCACTG
	15	GCTGAGAATG	AGGTTCT	TTTTGGAACG
	AACGGAACAA	AGGATGAGCA	GCCCGAGGGC	TCAGAG
	CTCAGCTCCCACT	GGACACCCTT	CACGGTTCCC	AAGGCCAGCC
	AGCTGCAGCA	GTGACGCCTG	GAAGGACATC	TGGTGGTCCT
	TAGGGGAGTG	GCCCCTCCTG	AGCCCTGCGA	GCAGCGTCCT
	TTTCTCTTTC	CCTCAGGCAG	CGGCTGTGTG	20
	AACCGCTGGC	TGCTGTTGTG	CCTCATCTCT	GGGCACATTG
	CCTGCTTCCC	CCCAGCGCCG	GCTTCTCTCC	TCAGAGCGCC
	TGCTACTCCA	TCCCCGGCAG	GGAGGGACCG	TCAGCTCACA
	AGGCCCTCTT	TGTTTCTCTG	TCCCAGACAT	AAGCCCAAGG
	GGCCCCTGCA	CCCAAGGGAC	CCTGTCCCTC	GGTGGCCTCC
	CCAGGCCCTT	25	GGACACGACA	GTTCTCTCTCA
	GGCAGGTGGG	CTTTGTGGTC	CTCGCCGCC	CTGGCCACAT
	CGCCCTCTCC	TCTTACACCT	GGTACCTTC	GAATGTTTCA
	GAGCGCAGGG	CCGTTCTCCC	TCGTGTCTC	TGGACCCACC
	CGCCCCCTTC	TGCCCTGTTT	GCGCAGGGAC	ATCACCCACA
	TGCCCCAGCT	CTCGGACCCT	30	GCAGCTCTGT
	GTCCCAGGCC	ACAGCAAAGG	TCTGTTGAAC	CCCTCCCTCC
	ATTCCCAGTT	ATCTGGGTCC	TCTGGATTCT	TCTGTTTCTT

GAATCAGGCT CTGCTTTCCC CCTAGCCACT ACAGGCAGCC  
 TCTGACAGTG CCGCTTTACT TGCATTCTGC AGCAATTACA TGTGTCTTTT  
 TGATCCTTGC CCAACTTCCC TCCCTCTCCC AGCTCCTGGC CCCTGGCCCA  
 GGGCCCTCT TGCTGTTTTT ACCTTTGTTC CTTGGGGCCT AGTACCCAGC  
 5 AAGCACCCAA ATGGGGGAGG TTTTGGGATG AGAGGAGGAA  
 ACGTGTATAC CTGTAACATC TGGTGGCTCT TCCCCAGAA GTTTGTGTTC  
 ATACATAATT GTTTTCCACG CTGGATCATA ATGTGACGTG CAGTTCTGCC  
 CTGTGCTGGG GAGCCACATG AAGCTTCCCC TGGCTAACTT  
 GCTACCCCGC AGCAATCCCA GTGTGGCCGT CTGCTTGCTA  
 10 AAAAATGGAT CTGTGCTCAT CTGTAAAAA AAAAAAAAAA AAAAAA

SEQ ID NO: 5 – PCa3\_splice variant protein

15 QGEGMRMAATAWAGLQGPPLPTLCPAVRTGLYCRDQA  
 HAERWAMTSETSSSGSHCARMLRRRAQEEDSTVLIDVS  
 PPEAEKRGSYGSTAHASEPPGGQAAACRAGSPAKPRI  
 DFLVWEEEDLKLD RQQDSAARDRTDMHRTWRETFDND  
 20 LRAAGLCVDQDQDVGDNNTTVHYALLSASWAVLCYYAE  
 DLRLKLPLQELPNQASNWSAGLLAWLGIPNVLLLEVVPD  
 VPPEYYSCRFRVNLKPRFLGSDNQDTFFTSTKRHQILF  
 EILAKTPYGHEKKNLLGIHQLLAEGVLSAAFPLHDGPFK  
 TPPEGPQAPRLNQRQVLFQHWARWGKWNKYQPLDHV  
 25 RRYFGEKVALYFAWLGFYTGWLLPAAVVGTLVFLVGCF  
 LVFSDIPTQELCGSKDSFEMCPLCLDCPFWLLSSACAL  
 AQAGRLFDHGGTVFFSLFMALWAVLLLEYWKRKSATLA  
 YRWDCSDYEDTEERPRPQFAASAPMTAPNPITGEDEP  
 YFPERSRARRMLAGSVVIVVMVAVVVMCLVSIILYRAIM  
 30 AIVVSRSGNTLLAAWASRIASLTGSVVNLVFIILSKIYV  
 SLAHVLTRWEMHRTQTKFEDAFTLKVFIFQFVNIFYSSP

VYIAFFKGRFVGYPGNYHTLFGVRNEECAAGGCLIELA  
QELLVIMVGKQVINNMQEVLPKLKGWWQKFRLLRSKKR  
KAGASAGASQGPWEDDYELVPCEGLFDEYLEMVLQFG  
FVTIFVAACPLAPLFPALLNNWVEIRLDARKFVCEYRRPV  
5 AERAQDIGIWFHILAGLTHLAVISNAFLAFSSDFLPRA  
YYRWTRAHDLRGFLNFTLARAPSSFAAAHNRTCRYRA  
FRDDDGHYSQTYWNLLAIRLAFVIVFEVAEAPAGSPIH  
GMRPRPCALPNSSTW

10

The cDNAs of the invention (SEQ ID NO: 2 or 4 from prostate-specific mRNA) are useful for screening for disseminated tumor cells by PCR, and immunotherapy of prostate cancer using transfected antigen-presenting cells. Proteins of the invention (SEQ ID NO: 3 or 5) are useful as a prostate specific  
15 antigen, for screening for disseminated tumor cells by antibody, and for immunotherapy of prostate cancer using protein or related peptides for induction of CD4+ cells and CD8+ cells.

Sequences represent new prostate-specific transcripts that are expressed in virtually all prostate cancers. Fragments of these sequences have been  
20 described as being prostate-specific by using a mere in silico approach (see Walker, M. G. et al., Prediction of gene function by genome-scale expression analysis: prostate cancer-associated genes. *Genome Res.* 9: 1198-1203 (1999)). However, in accordance with the present invention, these sequences were detected in a GX2000 database search and extended. Experimental data for their  
25 tissue specificity was demonstrated.

In a GX2000 screen for Genes specifically expressed in prostate tissue the EST AF109300 (SEQ ID NO: 1) was identified. Electronic northern provided evidence for expression in about 50% of all normal and malignant prostate tissue samples in the GX2000 database. Background expression in all available normal  
30 tissues was undetectable. AF109300 is part of a known cDNA library (Incyte

Pharmaceuticals) and is called D-PCa-3 (prostate cancer associated protein). The prostate-cancer association was determined by an *in silico* method called "guilt-by-association" (see Walker et al, *supra*).

5 The sequence was also identified in a GX2000 screen and provided experimental evidence for prostate-specific expression by hybridization of a multiple tissue expression array (from BD Clontech) with a radioactive probe representing AF109300. Prostate-specific expression was also demonstrated by quantitative PCR using a human cDNA panel (from BD Clontech) containing cDNA of 16 major organs as templates. Moreover, the sequence was extended  
10 using 5'- and 3'-RACE (rapid amplification of cDNA ends). The subsequently identified open reading frame comprises nucleotides coding for a putative protein containing 7 predicted transmembrane domains and a truncated DUF590 domain. A secreting signal is lacking. However, the ORF is still open at the 5'-end, indicating that the sequence is incomplete at its 5'-end. The present  
15 application also discloses a splice variant.

The sequences disclosed herein incorporate the polynucleotide transcript of SEQ ID NO: 2 and 4 and the derived amino acid sequences (SEQ ID NO: 2 or 4 and 5) from said transcripts present targets for chemotherapeutic agents,  
20 especially anti-cancer agents, including antibodies specific for such polypeptides.

The cancer-related polynucleotide sequences disclosed herein correspond to gene sequences whose expression is indicative of the cancerous status of a given cell. Such sequences are substantially identical to SEQ ID NO: 2 and 4,  
25 which represent transcripts identified from the GenBank EST database and which exhibit cancer-specific expression. The polynucleotides of the invention are those that correspond to the sequence of SEQ ID NO: 2 or 4.

The nucleotides and polypeptides, as gene products, used in the processes  
30 of the present invention may comprise a recombinant polynucleotide or polypeptide,

a natural polynucleotide or polypeptide, or a synthetic polynucleotide or polypeptide, or a recombinant polynucleotide or polypeptide.

Fragments of such polynucleotides and polypeptides as are disclosed herein  
5 may also be useful in practicing the processes of the present invention. For example, a fragment, derivative or analog of the polypeptide (SEQ ID NO: 3 or 5) may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one  
10 encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory  
15 sequence or a sequence which is employed for purification of the mature polypeptide (such as a histidine hexapeptide) or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

20 In another aspect, the present invention relates to an isolated polypeptide, including a purified polypeptide, comprising an amino acid sequence at least 90% identical to the amino acid sequence of SEQ ID NO: 3 or 5. In preferred embodiments, said isolated polypeptide comprises an amino acid sequence having sequence identity of at least 95%, preferably at least about 98%, and especially is  
25 identical to, the sequence of SEQ ID NO: 3 or 5. The present invention also includes isolated active fragments of such polypeptides where said fragments retain the biological activity of the polypeptide or where such active fragments are useful as specific targets for cancer treatment, prevention or diagnosis. Thus, the present invention relates to any polypeptides, or fragments thereof, with sufficient sequence  
30 homology to the sequences disclosed herein as to be useful in the production of antibodies that react with (i.e., are selective or specific for) the polypeptide of SEQ



ID NO: 3 or 5 so as to be useful in targeting cells that exhibit such polypeptides, or fragments, on their surfaces, thereby providing targets for such antibodies and therapeutic agents associated with such antibodies.

5 The polynucleotides and polypeptides useful in practicing the processes of the present invention may likewise be obtained in an isolated or purified form. In addition, the polypeptide disclosed herein as being useful in practicing the processes of the invention are believed to be surface proteins present on cells, such as cancerous cells. Precisely how such cancer-linked proteins are used in the processes of the invention may thus differ depending on the therapeutic approach  
10 used. For example, cell-surface proteins, such as receptors, are desirable targets for cytotoxic antibodies that can be generated against the polypeptides disclosed herein.

The present invention relates to an isolated polypeptide, encoded by one of  
15 the polynucleotide transcripts disclosed herein, comprising an amino acid sequence homologous to the amino acid sequence of SEQ ID NO: 3 or 5, wherein any difference between amino acid sequence in the isolated polypeptide and the sequence of SEQ ID NO: 3 or 5 is due solely to conservative amino acid substitutions and wherein said isolated polypeptide comprises at least one  
20 immunogenic fragment. In a preferred embodiment, the present invention encompasses an isolated polypeptide comprising the amino acid sequence of SEQ ID NO: 3 or 5.

Methods of producing recombinant cells and vectors useful in preparing  
25 the polynucleotides and polypeptides disclosed herein are well known to those skilled in the molecular biology art. See, for example, Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), Wu et al., *Methods in Gene Biotechnology* (CRC Press, New York, NY, 1997), and *Recombinant Gene Expression Protocols*, in *Methods in Molecular Biology*,  
30 Vol. 62, (Tuan, ed., Humana Press, Totowa, NJ, 1997), the disclosures of which are hereby incorporated by reference.

In one aspect, the present invention relates to a process for identifying an agent that modulates the activity of a cancer-related gene comprising:

(a) contacting a compound with a cell containing a gene that corresponds to a polynucleotide having the sequence of SEQ ID NO: 2 or 4 and under  
5 conditions promoting the expression of said gene; and

(b) detecting a difference in expression of said gene relative to when said compound is not present

thereby identifying an agent that modulates the activity of a cancer-related gene.

10

In specific embodiments of such process the cell is a cancer cell and the difference in expression is a decrease in expression. Such polynucleotides may also include those that have the sequence of SEQ ID NO: 2 or 4.

15

In another aspect, the present invention relates to a process for identifying an anti-neoplastic agent comprising contacting a cell exhibiting neoplastic activity with a compound first identified as a cancer related gene modulator using an assay process disclosed herein and detecting a decrease in said neoplastic activity after said contacting compared to when said contacting does not occur.

20

Such neoplastic activity may include accelerated cellular replication and/or metastasis, and the decrease in neoplastic activity preferably results from the death of the cell.

25

The present invention also relates to a process for identifying an anti-neoplastic agent comprising administering to an animal exhibiting a cancer condition an effective amount of an agent first identified according to a process of one of one of the assays disclosed according to the invention and detecting a decrease in said cancerous condition.

30

In specific embodiments of the present invention, the genes useful for the invention comprise genes that correspond to the polynucleotide having the

sequence of SEQ ID NO: 2 or 4, or may comprise the sequence of the cDNA (SEQ ID NO: 2 or 4) disclosed herein.

In accordance with the present invention, such assays rely on methods of determining the activity of the gene in question. Such assays are advantageously  
5 based on model cellular systems using cancer cell lines, primary cancer cells, or cancerous tissue samples that are maintained in growth medium and treated with compounds at a single concentration or at a range of concentrations. At specific times after treatment, cellular RNAs are conveniently isolated from the treated cells or tissues, which RNAs are indicative of expression of selected genes. The  
10 cellular RNA is then divided and subjected to differential analysis that detects the presence and/or quantity of specific RNA transcripts, which transcripts may then be amplified for detection purposes using standard methodologies, such as, for example, reverse transcriptase polymerase chain reaction (RT-PCR), etc. The presence or absence, or concentration levels, of specific RNA transcripts are  
15 determined from these measurements. The polynucleotide sequences disclosed herein are readily used as probes for the detection of such RNA transcripts and thus the measurement of gene activity and expression.

The polynucleotides of the invention can include fully operational genes  
20 with attendant control or regulatory sequences or merely a polynucleotide sequence encoding the corresponding polypeptide or an active fragment or analog thereof.

Because expression of the polynucleotide sequences disclosed herein are  
25 specific to the cancerous state, useful gene modulation is downward modulation, so that, as a result of exposure to an anti-neoplastic agent identified by the screening assays herein, the corresponding gene of the cancerous cell is expressed at a lower level (or not expressed at all) when exposed to the agent as compared to the expression when not exposed to the agent. For example, the  
30 gene sequence disclosed herein (SEQ ID NO: 2 or 4) correspond to a gene expressed at a higher level in cells of prostate cancer than in normal prostate

cells. Thus, where said chemical agent causes this gene of the tested cell to be expressed at a lower level than the same genes of the reference, this is indicative of downward modulation and indicates that the chemical agent to be tested has anti-neoplastic activity.

5

In carrying out the assays disclosed herein, relative anti-neoplastic activity may be ascertained by the extent to which a given chemical agent modulates the expression of genes present in a cancerous cell. Thus, a first chemical agent that modulates the expression of a gene associated with the cancerous state (i.e., a  
10 gene corresponding to the polynucleotide transcript of SEQ ID NO: 2 or 4) to a larger degree than a second chemical agent tested by the assays of the invention is thereby deemed to have higher, or more desirable, or more advantageous, anti-neoplastic activity than said second chemical agent.

15 The gene expression to be measured is commonly assayed using RNA expression as an indicator. Thus, the greater the level of RNA (for example, messenger RNA or mRNA) detected the higher the level of expression of the corresponding gene. Thus, gene expression, either absolute or relative, is determined by the relative expression of the RNAs encoded by such genes.

20

RNA may be isolated from samples in a variety of ways, including lysis and denaturation with a phenolic solution containing a chaotropic agent (e.g., trizol) followed by isopropanol precipitation, ethanol wash, and resuspension in aqueous solution; or lysis and denaturation followed by isolation on solid support, such as a  
25 Qiagen resin and reconstitution in aqueous solution; or lysis and denaturation in non-phenolic, aqueous solutions followed by enzymatic conversion of RNA to DNA template copies.

30 Normally, prior to applying the processes of the invention, steady state RNA expression levels for the genes, and sets of genes, disclosed herein will have been obtained. It is the steady state level of such expression that is affected by potential

anti-neoplastic agents as determined herein. Such steady state levels of expression are easily determined by any methods that are sensitive, specific and accurate. Such methods include, but are in no way limited to, real time quantitative polymerase chain reaction (PCR), for example, using a Perkin-Elmer 7700  
5 sequence detection system with gene specific primer probe combinations as designed using any of several commercially available software packages, such as Primer Express software., solid support based hybridization array technology using appropriate internal controls for quantitation, including filter, bead, or microchip based arrays, solid support based hybridization arrays using, for example,  
10 chemiluminescent, fluorescent, or electrochemical reaction based detection systems.

The gene expression indicative of a cancerous state need not be characteristic of every cell of a given tissue. Thus, the methods disclosed herein  
15 are useful for detecting the presence of a cancerous condition within a tissue where less than all cells exhibit the complete pattern. Thus, for example, a selected gene corresponding to the sequence of SEQ ID NO: 2 or 4, may be found, using appropriate probes, either DNA or RNA, to be present in as little as 60% of cells derived from a sample of tumorous, or malignant, tissue. In a highly  
20 preferred embodiment, such gene pattern is found to be present in at least 100% of cells drawn from a cancerous tissue and absent from at least 100% of a corresponding normal, non-cancerous, tissue sample.

Expression of a gene may be related to copy number, and changes in  
25 expression may be measured by determining copy number. Such change in gene copy number may be determined by determining a change in expression of messenger RNA encoded by a particular gene sequence, especially that of SEQ ID NO: 2 or 4. Also in accordance with the present invention, said gene may be a cancer initiating or facilitating gene. In carrying out the methods of the present  
30 invention, a cancer facilitating gene is a gene that, while not directly initiating tumor formation or growth, acts, such as through the actions of its expression

product, to direct, enhance, or otherwise facilitate the progress of the cancerous condition, including where such gene acts against genes, or gene expression products, that would otherwise have the effect of decreasing tumor formation and/or growth.

5

Although the expression of a gene corresponding to the sequence of SEQ ID NO: 2 or 4 may be indicative of a cancerous status for a given cell, the mere presence of such gene may not alone be sufficient to achieve a malignant condition and thus the level of expression of such gene may also be a significant factor in determining the attainment of a cancerous state. Thus, it becomes essential to also determine the level of expression of a gene as disclosed herein, including substantially similar sequences, as a separate means of diagnosing the presence of a cancerous status for a given cell, groups of cells, or tissues, either in culture or *in situ*.

15

The level of expression of the polypeptide disclosed herein is also a measure of gene expression, such as the polypeptide having sequence identical, or similar to, SEQ ID NO: 3 or 5.

20

In accordance with the foregoing, the present invention specifically contemplates a method for determining the cancerous status of a cell to be tested, comprising determining the level of expression in said cell of a gene that includes the nucleotide sequence of SEQ ID NO: 2 or 4, including sequences substantially identical to said sequence, or characteristic fragments thereof, or the complements of any of these and then comparing said expression to that of a cell known to be non-cancerous whereby the difference in said expression indicates that said cell to be tested is cancerous.

25

In accordance with the invention, although gene expression for a gene that includes as a portion thereof the sequence of SEQ ID NO: 2 or 4, is preferably determined by use of a probe that is a fragment of such nucleotide

30

sequence, it is to be understood that the probe may be formed from a different portion of the gene. Expression of the gene may be determined by use of a nucleotide probe that hybridizes to messenger RNA (mRNA) transcribed from a portion of the gene other than the specific nucleotide sequence disclosed herein.

5

It should be noted that there are a variety of different contexts in which genes have been evaluated as being involved in the cancerous process. Thus, some genes may be oncogenes and encode proteins that are directly involved in the cancerous process and thereby promote the occurrence of cancer in an animal. In addition, other genes may serve to suppress the cancerous state in a given cell or cell type and thereby work against a cancerous condition forming in an animal. Other genes may simply be involved either directly or indirectly in the cancerous process or condition and may serve in an ancillary capacity with respect to the cancerous state. All such types of genes are deemed with those to be determined in accordance with the invention as disclosed herein. Thus, the gene determined by said process of the invention may be an oncogene, or the gene determined by said process may be a cancer facilitating gene, the latter including a gene that directly or indirectly affects the cancerous process, either in the promotion of a cancerous condition or in facilitating the progress of cancerous growth or otherwise modulating the growth of cancer cells, either *in vivo* or *ex vivo*. In addition, the gene determined by said process may be a cancer suppressor gene, which gene works either directly or indirectly to suppress the initiation or progress of a cancerous condition. Such genes may work indirectly where their expression alters the activity of some other gene or gene expression product that is itself directly involved in initiating or facilitating the progress of a cancerous condition. For example, a gene that encodes a polypeptide, either wild or mutant in type, which polypeptide acts to suppress of tumor suppressor gene, or its expression product, will thereby act indirectly to promote tumor growth.

30

As noted previously, polynucleotides encoding the same protein as SEQ ID NO: 2 or 4, regardless of the percent identity of such sequences, are also specifically contemplated by any of the methods of the present invention that rely on any or all of said sequences, regardless of how they are otherwise described or limited. Thus, any such sequences are available for use in carrying out any of the methods disclosed according to the invention. Such sequences also include any open reading frames, as defined herein, present within the sequence of SEQ ID NO: 2 or 4.

Because a gene disclosed according to the invention "corresponds to" a polynucleotide having the sequence of SEQ ID NO: 2 or 4, said gene encodes an RNA (processed or unprocessed, including naturally occurring splice variants and alleles) that is at least 90% identical, preferably at least 95% identical, most preferably at least 98% identical to, and especially identical to, an RNA that would be encoded by, or be complementary to, such as by hybridization with, a polynucleotide having the indicated sequence. In addition, genes including sequences at least 90% identical to the sequence of SEQ ID NO: 2 or 4, preferably at least about 95% identical to such a sequence, more preferably at least about 98% identical to such sequence and most preferably comprising such sequence are specifically contemplated by all of the processes of the present invention. Sequences encoding the same proteins as any of these sequences, regardless of the percent identity of such sequences, are also specifically contemplated by any of the methods of the present invention that rely on any or all of said sequences, regardless of how they are otherwise described or limited. The polynucleotide sequences of the invention also include any open reading frames, as defined herein, present within the sequence of SEQ ID NO: 2 or 4.

The sequences disclosed herein may be genomic in nature and thus represent the sequence of an actual gene, such as a human gene, or may be a cDNA sequence derived from a messenger RNA (mRNA) and thus represent contiguous exonic sequences derived from a corresponding genomic sequence,



or they may be wholly synthetic in origin for purposes of practicing the processes of the invention. Because of the processing that may take place in transforming the initial RNA transcript into the final mRNA, the sequences disclosed herein may represent less than the full genomic sequence. They may also represent sequences derived from ribosomal and transfer RNAs. Consequently, the gene as present in the cell (and representing the genomic sequence) and the polynucleotide transcripts disclosed herein, including cDNA sequences, may be identical or may be such that the cDNAs contain less than the full genomic sequence. Such genes and cDNA sequences are still considered "corresponding sequences" (as defined elsewhere herein) because they both encode the same or related RNA sequences (i.e., related in the sense of being splice variants or RNAs at different stages of processing). Thus, by way of non-limiting example only, a gene that encodes an RNA transcript, which is then processed into a shorter mRNA, is deemed to encode both such RNAs and therefore encodes an RNA complementary to (using the usual Watson-Crick complementarity rules), or that would otherwise be encoded by, a cDNA (for example, a sequence as disclosed herein). Thus, the sequences disclosed herein correspond to genes contained in the cancerous cells (here, prostate cancer) and are used to determine gene activity or expression because they represent the same sequence or are complementary to RNAs encoded by the gene. Such a gene also includes different alleles and splice variants that may occur in the cells used in the methods of the invention, such as where recombinant cells are used to assay for anti-neoplastic agents and such cells have been engineered to express a polynucleotide as disclosed herein, including cells that have been engineered to express such polynucleotides at a higher level than is found in non-engineered cancerous cells or where such recombinant cells express such polynucleotides only after having been engineered to do so. Such engineering includes genetic engineering, such as where one or more of the polynucleotides disclosed herein has been inserted into the genome of such cell or is present in a vector.

Such cells, especially mammalian cells, may also be engineered to express on their surfaces a polypeptide of the invention for testing with antibodies or other agents capable of masking such polypeptides and thereby removing the cancerous nature of the cell. Such engineering includes both  
5 genetic engineering, where the genetic complement of the cells is engineered to express the polypeptide, as well as non-genetic engineering, whereby the cell has been physically manipulated to incorporate a polypeptide of the invention in its plasma membrane, such as by direct insertion using chemical and/or other agents to achieve this result.

10 In accordance with the foregoing, the present invention includes anti-cancer agents that are themselves either polypeptides, or small chemical entities, that affect the cancerous process, including initiation, suppression or facilitation of tumor growth, either *in vivo* or *ex vivo*. Said cancer modulating agent will have  
15 the effect of decreasing gene expression.

The present invention thus also relates to a method for treating cancer comprising contacting a cancerous cell with an agent having activity against an expression product encoded by a gene or polynucleotide sequence as disclosed  
20 herein, such as one having, or corresponding to, the nucleotide sequence of SEQ ID NO: 2 or 4. The present invention also relates to a process for treating cancer comprising contacting a cancerous cell with an agent having activity against an expression product encoded by a gene or polynucleotide sequence corresponding to SEQ ID NO: 2 or 4. In one such embodiment, the cancerous  
25 cell is contacted *in vivo*. In another such embodiment, said agent has affinity for said expression product. In a preferred embodiment, such agent is an antibody disclosed herein, such as an antibody that is specific or selective for, or otherwise reacts with, a polypeptide of the invention. In a preferred embodiment, the expression product is a polypeptide incorporating the amino acid sequence of  
30 SEQ ID NO: 3 or 5.

The present invention is also directed to such uses of the compositions of polypeptides and antibodies disclosed herein. Such uses include a process for treating cancer in an animal afflicted therewith comprising administering to said animal an amount of an immunogenic composition of one or more of the polypeptides disclosed herein where such amount is an amount sufficient to elicit the production of cytotoxic T lymphocytes specific for a polypeptide of the invention, preferably a polypeptide incorporating a sequence of SEQ ID NO: 3 or 5. In a preferred embodiment, the animal to be so treated is a human patient.

10 The proteins encoded by the genes disclosed herein due to their expression, or elevated expression, in cancer cells, represent highly useful therapeutic targets for "targeted therapies" utilizing such affinity structures as, for example, antibodies coupled to some cytotoxic agent. In such methodology, it is advantageous that nothing need be known about the endogenous ligands or  
15 binding partners for such cell surface molecules. Rather, an antibody or equivalent molecule that can specifically recognize the cell surface molecule (which could include an artificial peptide, a surrogate ligand, and the like) that is coupled to some agent that can induce cell death or a block in cell cycling offers therapeutic promise against these proteins. Thus, such approaches include the  
20 use of so-called suicide "bullets" against intracellular proteins. For example, monoclonal antibodies may readily be produced by methods well known in the art, for example, the method of Kohler and Milstein (see: *Nature*, **256**:495 (1975).

25 With the advent of methods of molecular biology and recombinant technology, it is now possible to produce antibody molecules by recombinant means and thereby generate gene sequences that code for specific amino acid sequences found in the polypeptide structure of the antibodies. Such antibodies can be produced by either cloning the gene sequences encoding the polypeptide chains of said antibodies or by direct synthesis of said polypeptide chains, with  
30 *in vitro* assembly of the synthesized chains to form active tetrameric (H<sub>2</sub>L<sub>2</sub>) structures with affinity for specific epitopes and antigenic determinants. This has

permitted the ready production of antibodies having sequences characteristic of neutralizing antibodies from different species and sources.

Regardless of the source of the antibodies, or how they are recombinantly constructed, or how they are synthesized, *in vitro* or *in vivo*, using transgenic animals, such as cows, goats and sheep, using large cell cultures of laboratory or commercial size, in bioreactors or by direct chemical synthesis employing no living organisms at any stage of the process, all antibodies have a similar overall 3 dimensional structure. This structure is often given as  $H_2L_2$  and refers to the fact that antibodies commonly comprise 2 light (L) amino acid chains and 2 heavy (H) amino acid chains. Both chains have regions capable of interacting with a structurally complementary antigenic target. The regions interacting with the target are referred to as "variable" or "V" regions and are characterized by differences in amino acid sequence from antibodies of different antigenic specificity.

The variable regions of either H or L chains contains the amino acid sequences capable of specifically binding to antigenic targets. Within these sequences are smaller sequences dubbed "hypervariable" because of their extreme variability between antibodies of differing specificity. Such hypervariable regions are also referred to as "complementarity determining regions" or "CDR" regions. These CDR regions account for the basic specificity of the antibody for a particular antigenic determinant structure.

The CDRs represent non-contiguous stretches of amino acids within the variable regions but, regardless of species, the positional locations of these critical amino acid sequences within the variable heavy and light chain regions have been found to have similar locations within the amino acid sequences of the variable chains. The variable heavy and light chains of all antibodies each have 3 CDR regions, each non-contiguous with the others (termed L1, L2, L3, H1, H2,

H3) for the respective light (L) and heavy (H) chains. The accepted CDR regions have been described by Kabat et al., *J. Biol. Chem.* 252:6609-6616 (1977).

5 In all mammalian species, antibody polypeptides contain constant (i.e., highly conserved) and variable regions, and, within the latter, there are the CDRs and the so-called "framework regions" made up of amino acid sequences within the variable region of the heavy or light chain but outside the CDRs.

10 The antibodies disclosed according to the invention may also be wholly synthetic, wherein the polypeptide chains of the antibodies are synthesized and, possibly, optimized for binding to the polypeptides disclosed herein as being receptors. Such antibodies may be chimeric or humanized antibodies and may be fully tetrameric in structure, or may be dimeric and comprise only a single heavy and a single light chain. Such antibodies may also include fragments, such as Fab and F(ab<sub>2</sub>)' fragments, capable of reacting with and binding to any of the  
15 polypeptides disclosed herein as being receptors.

In one aspect, the present invention relates to immunoglobulins, or antibodies, as described herein, that react with, especially where they are  
20 specific for, the polypeptides having amino acid sequences as disclosed herein, preferably those having the amino acid sequence of SEQ ID NO: 3 or 5. Such antibodies may commonly be in the form of a composition, especially a pharmaceutical composition. Such antibodies, by themselves, may have therapeutic value in that they are able to bind to, and thereby tie up, surface sites  
25 on cancerous cells. Where such sites have some type of function to perform (i.e., where they are surface enzymes, or channel structures, or structures that otherwise facilitate, actively or passively, the transport of nutrients and other vital materials to the cell. Such nutrients serve to facilitate the growth and replication of the cell and molecules that bind to such sites and thereby interfere with such  
30 activities can prove to have a therapeutic effect in that the result of such binding is to remove sources of nutrients from such cells, thereby interfering with growth

and replication. In like manner, such binding may serve to remove vital enzyme activities from the cell's functional repertoire, thereby also interfering with viability and/or the ability of the cell to multiply or metastasize. In addition, by binding to such surface sites, the antibodies may serve to prevent the cells from reacting to environmental agents, such as cytokines and the like, that may facilitate growth, replication and metastasis, thereby further reducing the cancerous status of such cell and ameliorating the cancerous condition in a patient, even without proving fatal to the cell or cells so affected.

The methods of the present invention also include processes wherein the cancer cell is contacted *in vivo* as well as *ex vivo* with an agent that comprises a portion, or is part of an overall molecular structure, having affinity for an expression product of a gene corresponding to a polynucleotide sequence as disclosed herein, preferably where the expression product is a cell surface structure, most preferably a polypeptide as disclosed herein, such as one that comprises an amino acid sequence homologous to, or including, SEQ ID NO: 3 or 5. In one such embodiment, said portion having affinity for said expression product is an antibody, especially where said expression product is a polypeptide or oligopeptide or comprises an oligopeptide portion, or comprises a polypeptide.

In another aspect, the present invention also relates to an antibody that reacts with a polypeptide as disclosed herein, preferably a polypeptide comprising the amino acid sequence of SEQ ID NO: 3 or 5. Such an antibody may be polyclonal, monoclonal, recombinant or synthetic in origin. In one such embodiment, said antibody is associated, either covalently or non-covalently, with a cytotoxic agent, for example, an apoptotic agent. It is thus contemplated that the antibody acts a targeted vector for guiding an associated therapeutic agent to a cancerous cell, such as a cell expressing a polypeptide homologous to, if not identical to, a polypeptide as disclosed herein.

Where the cytotoxic agent is itself a polypeptide, said may be linked directly to an antibody specific for a surface target on a cancer cell, such as where the polypeptide represents an extension of the amino acid chain of the antibody. In alternative embodiments, such molecules may be covalently linked  
5 through a linker sequence of long or short duration, such as an amino acid sequence of 5 to 10 residues in length. Where the cytotoxic agents is some small organic molecule, such as a small organic compound, or some type of apoptotic agent, this may be covalently bonded to the antibody molecule or may be attached by some other type of non-covalent linkage, including hydrophobic and  
10 electrostatic linkages. Methods for forming such linkages, especially covalent linkages, are well known to those skilled in the art.

The antibodies disclosed herein may also serve as targeting vectors for much larger structures, such as liposomes. In one such embodiment, an antibody  
15 is part of, or otherwise linked to, or associated with, a membranous structure, preferably a liposome or possibly some type of cellular organelle, which acts as a reservoir for a cytotoxic agent, such as ricin. The antibody then acts to target said liposome to a cancerous tissue in an animal, whereupon the liposome provides a source of cytotoxic agents for localized treatment of a solid tumor or other type of  
20 neoplasm.

The present invention further encompasses an immunogenic composition comprising a polypeptide disclosed herein, as well as compositions formed using antibodies specific for these polypeptides.

25

Methods well known in the art for making formulations are found in, for example, *Remington: The Science and Practice of Pharmacy*, (19th ed.) Ed. A.R. Gennaro, 1995, Mack Publishing Company, Easton, PA. Formulations for parenteral administration may, for example, contain excipients, sterile water, or  
30 saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes. Biocompatible, biodegradable lactide polymer,

lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of the compounds. Other potentially useful parenteral delivery systems for agonists of the invention include ethylenevinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain excipients, or example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel. It should be noted that, where the therapeutic agent to be administered is an immunoconjugate, these sometimes contain chemical linkages that are somewhat labile in aqueous media and therefor must be stored prior to administration in a more stable environment, such as in the form of a lyophilized powder.

Such an agent can be a single molecular structure, comprising both affinity portion and anti-cancer activity portions, wherein said portions are derived from separate molecules, or molecular structures, possessing such activity when separated and wherein such agent has been formed by combining said portions into one larger molecular structure, such as where said portions are combined into the form of an adduct. Said anti-cancer and affinity portions may be joined covalently, such as in the form of a single polypeptide, or polypeptide-like, structure or may be joined non-covalently, such as by hydrophobic or electrostatic interactions, such structures having been formed by means well known in the chemical arts. Alternatively, the anti-cancer and affinity portions may be formed from separate domains of a single molecule that exhibits, as part of the same chemical structure, more than one activity wherein one of the activities is against cancer cells, or tumor formation or growth, and the other activity is affinity for an expression product produced by expression of genes related to the cancerous process or condition.

In one embodiment of the present invention, a chemical agent, such as a protein or other polypeptide, is joined to an agent, such as an antibody, having



affinity for an expression product of a cancerous cell, such as a polypeptide or protein encoded by a gene related to the cancerous process, preferably a gene as disclosed herein according to the present invention, most preferably a polypeptide sequence disclosed herein. Thus, where the presence of said  
5 expression product is essential to tumor initiation and/or growth, binding of said agent to said expression product will have the effect of negating said tumor promoting activity. In one such embodiment, said agent is an apoptosis-inducing agent that induces cell suicide, thereby killing the cancer cell and halting tumor growth.

10

Other genes within the cancer cell that are regulated in a manner similar to that of the genes disclosed herein and thus change their expression in a coordinated way in response to chemical compounds represent genes that are located within a common metabolic, signaling, physiological, or functional  
15 pathway so that by analyzing and identifying such commonly regulated groups of genes (groups that include the gene, or similar sequences, disclosed according to the invention, one can (a) assign known genes and novel genes to specific pathways and (b) identify specific functions and functional roles for novel genes that are grouped into pathways with genes for which their functions are already  
20 characterized or described. For example, one might identify a group of 10 genes, at least one of which is the gene as disclosed herein, that change expression in a coordinated fashion and for which the function of one, such as the polypeptide encoded by the sequence disclosed herein, is known then the other genes are thereby implicated in a similar function or pathway and may thus play a role in  
25 the cancer-initiating or cancer-facilitating process. In the same way, if a gene were found in normal cells but not in cancer cells, or happens to be expressed at a higher level in normal as opposed to cancer cells, then a similar conclusion may be drawn as to its involvement in cancer, or other diseases. Therefore, the processes disclosed according to the present invention at once provide a novel  
30 means of assigning function to genes, i.e. a novel method of functional genomics, and a means for identifying chemical compounds that have potential

therapeutic effects on specific cellular pathways. Such chemical compounds may have therapeutic relevance to a variety of diseases outside of cancer as well, in cases where such diseases are known or are demonstrated to involve the specific cellular pathway that is affected.

5

The polypeptides disclosed herein, preferably that of SEQ ID NO: 3 or 5, also find use as vaccines in that, where the polypeptide represents a surface protein present on a cancer cell, such polypeptide may be administered to an animal, especially a human being, for purposes of activating cytotoxic T lymphocytes (CTLs) that will be specific for, and act to lyse, cancer cells in said animal. Where used as vaccines, such polypeptides are present in the form of a pharmaceutical composition. The present invention may also employ polypeptides that have the same, or similar, immunogenic character as the polypeptide of SEQ ID NO: 3 or 5 and thereby elicit the same, or similar, immunogenic response after administration to an animal, such as an animal at risk of developing cancer, or afflicted therewith. Thus, the polypeptides disclosed according to the invention will commonly find use as immunogenic compositions.

Expression of a gene corresponding to a polynucleotide disclosed herein, when in normal tissues, may indicate a predisposition towards development of prostate cancer. The encoded polypeptide might then present a potentially useful cell surface target for therapeutic molecules such as cytolytic antibodies, or antibodies attached to cytotoxic, or cytolytic, agents.

The present invention specifically contemplates use of antibodies against the polypeptides encoded by the polynucleotides corresponding to the genes disclosed herein, whereby said antibodies are conjugates to one or more cytotoxic agents so that the antibodies serve to target the conjugated immunotoxins to a region of cancerous activity, such as a solid tumor. For many known cytotoxic agents, lack of selectivity has presented a drawback to their use as therapeutic agents in the treatment of malignancies. For example, the class of

two-chain toxins, consisting of a binding subunit (or B-chain) linked to a toxic subunit (A-chain) are extremely cytotoxic. Thus, such agents as ricin, a protein isolated from castor beans, kills cells at very low concentrations (even less than  $10^{-11}$  M) by inactivating ribosomes in said cells (see, for example, Lord et al.,

- 5 Ricin: structure, mode of action, and some current applications. *Faseb J*, **8**: 201-208 (1994), and Blättler et al., Realizing the full potential of immunotoxins. *Cancer Cells*, **1**: 50-55 (1989)). While isolated A-chains of protein toxins that functionally resemble ricin A-chain are only weakly cytotoxic for intact cells (in the concentration range of  $10^{-7}$  to  $10^{-6}$  M), they are very potent cytotoxic agents
- 10 inside the cells. Thus, a single molecule of the A-subunit of diphtheria toxin can kill a cell once inside (see: Yamaizumi et al., One molecule of diphtheria toxin fragment A introduced into a cell can kill the cell. *Cell*, **15**: 245-250, 1978).

The present invention solves this selectivity problem by using antibodies

15 specific for antigens present on cancer cells to target the cytotoxins to said cells. In addition, use of antibodies decreases toxicity because the antibodies are non-toxic until they reach the tumor and, because the cytotoxin is bound to the antibody, it is presented with less opportunity to cause damage to non-targeted tissues.

20

In addition, use of such antibodies alone can provide therapeutic effects on the tumor through the antibody-dependent cellular cytotoxic response (ADCC) and complement-mediated cell lysis mechanisms.

- 25 A number of recombinant immunotoxins (for example, consisting of Fv regions of cancer specific antibodies fused to truncated bacterial toxins) are well known (see, for example, Smyth et al., Specific targeting of chlorambucil to tumors with the use of monoclonal antibodies, *J. Natl. Cancer Inst.*, **76**(3):503-510 (1986); Cho et al., Single-chain Fv/folate conjugates mediate efficient lysis of
- 30 folate-receptor-positive tumor cells, *Bioconjug. Chem.*, **8**(3):338-346 (1997)). As noted in the literature, these may contain, for example, a truncated version of

*Pseudomonas* exotoxin as a toxic moiety but the toxin is modified in such a manner that by itself it does not bind to normal human cells, but it retains all other functions of cytotoxicity. Here, recombinant antibody fragments target the modified toxin to cancer cells which are killed, such as by direct inhibition of protein synthesis, or by concomitant induction of apoptosis. Cells that are not recognized by the antibody fragment, because they do not carry the cancer antigen, are not affected. Good activity and specificity has been observed for many recombinant immunotoxins in *in vitro* assays using cultured cancer cells as well as in animal tumor models. Ongoing clinical trials provide examples where the promising pre-clinical data correlate with successful results in experimental cancer therapy. (see, for example, Brinkmann U., Recombinant antibody fragments and immunotoxin fusions for cancer therapy, *In Vivo* (2000) 14:21-27).

While the safety of employing immunoconjugates in humans has been established, *in vivo* therapeutic results have been less impressive. Because clinical use of mouse MAbs in humans is limited by the development of a foreign anti-globulin immune response by the human host, genetically engineered chimeric human-mouse MAbs have been developed by replacing the mouse Fc region with the human constant region. In other cases, the mouse antibodies have been "humanized" by replacing the framework regions of variable domains of rodent antibodies by their human equivalents. Such humanized and engineered antibodies can even be structurally arranged to have specificities and effector functions determined by design and which characteristics do not appear in nature. The development of bispecific antibodies, having different binding ends so that more than one antigenic site can be bound, have proven useful in targeting cancer cells. Thus, such antibody specificity has been improved by chemical coupling to various agents such as bacterial or plant toxins, radionuclides or cytotoxic drugs and other agents. (see, for example, Bodey, B. et al). Genetically engineered monoclonal antibodies for direct anti-neoplastic treatment and cancer cell specific delivery of chemotherapeutic agents. *Curr Pharm Des* (2000) Feb;6(3):261-76). See also, Garnett, M. C., Targeted drug

conjugates: principles and progress. *Adv. Drug Deliv. Rev.* (2001 Dec 17) **53**(2):171-216; Brinkmann et al., Recombinant immunotoxins for cancer therapy. *Expert Opin Biol Ther.* (2001) **1**(4):693-702.

- 5           Among the cytotoxic agents specifically contemplated for use as immunoconjugates according to the present invention are Calicheamicin, a highly toxic enediyne antibiotic isolated from *Micromonospora echinospora ssp. Calichensis*, and which binds to the minor groove of DNA to induce double strand breaks and cell death (see: Lee et al., Calicheamicins, a novel
- 10 family of antitumor antibiotics. 1. Chemistry and partial structure of calicheamicin g<sub>1</sub>. *J Am Chem Soc*, **109**: 3464-3466 (1987); Zein et al., Calicheamicin gamma 11: an antitumor antibiotic that cleaves double-stranded DNA site specifically, *Science*, **240**: 1198-1201 (1988)). Useful derivatives of the calicheamicins include mylotarg and 138H11-Cam $\theta$ . Mylotarg is an immunoconjugate of a
- 15 humanized anti-CD33 antibody (CD33 being found in leukemic cells of most patients with acute myeloid leukemia) and N-acetyl gamma colicheamicin dimethyl hydrazide, the latter of which is readily coupled to an antibody of the present invention (in place of the anti-CD33 but which can also be humanized by substitution of human framework regions into the antibody during production as
- 20 described elsewhere herein) to form an immunoconjugate of the invention. (see: Hamann et al. Gemtuzumab Ozogamicin, A Potent and Selective Anti-CD33 Antibody-Calicheamicin Conjugate for Treatment of Acute Myeloid Leukemia, *Bioconjug. Chem.* **13**, 47-58 (2002)) For use with 138H11-Cam $\theta$ , 138H11 is an anti- $\gamma$ -glutamyl transferase antibody coupled to theta calicheamicin through a
- 25 disulfide linkage and found useful *in vitro* against cultured renal cell carcinoma cells. (see: Knoll et al., Targeted therapy of experimental renal cell carcinoma with a novel conjugate of monoclonal antibody 138H11 and calicheamicin  $\theta_1^1$ , *Cancer Res*, **60**: 6089-6094 (2000) The same linkage may be utilized to link this cytotoxic agent to an antibody of the present invention, thereby forming a
- 30 targeting structure for prostate cancer cells.

Also useful in forming the immunoconjugates of the invention is DC1, a disulfide-containing analog of adozelesin, that kills cells by binding to the minor groove of DNA, followed by alkylation of adenine bases. Adozelesin is a structural analog of CC-1065, an anti-tumor antibiotic isolated from microbial fermentation of *Streptomyces zelensis*, and is about 1,000 fold more toxic to cultured cell lines than other DNA interacting agents, such as cis-platin and doxorubicin. This agent is readily linked to antibodies through the disulfide bond of adozelesin. (see: Chari et al., Enhancement of the selectivity and antitumor efficacy of a CC-1065 analogue through immunoconjugate formation, *Cancer Res*, 55: 4079-4084 (1995)).

Maytansine, a highly cytotoxic microtubular inhibitor isolated from the shrub *Maytenus serrata* found to have little value in human clinical trials, is much more effective in its derivatized form, denoted DM1, containing a disulfide bond to facilitate linkage to antibodies, is up to 10-fold more cytotoxic (see: Chari et al., Immunoconjugates containing novel maytansinoids: promising anticancer drugs, *Cancer Res*, 52: 127-131 (1992)). These same *in vitro* studies showed that up to four DM1 molecules could be linked to a single immunoglobulin without destroying the binding affinity. Such conjugates have been used against prostate cancer antigens, such as the *neu*/HER2/*erbB*-2 antigen. (see: Goldmacher et al., Immunogen, Inc., (2002) *in press*); also see Liu, C. et al., Eradication of large colon tumor xenografts by targeted delivery of maytansinoids, *Proc. Natl. Acad. Sci. USA*, 93, 8618-8623 (1996)). For example, Liu et al. (1996) describes formation of an immunoconjugate of the maytansinoid cytotoxin DM1 and C242 antibody, a murine IgG1 immunoglobulin, available from Pharmacia and which has affinity for a mucin-like glycoprotein variably expressed by human colorectal cancers. The latter immunoconjugate was prepared according to Chari et al., *Cancer Res.*, 52:127-131 (1992) and was found to be highly cytotoxic against cultured colon cancer cells as well as showing anti-tumor effects *in vivo* in mice bearing subcutaneous COLO 205 human colon tumor xenografts using doses well below the maximum tolerated dose.

In addition, there are a variety of protein toxins (cytotoxic proteins), which include a number of different classes, such as those that inhibit protein synthesis: ribosome-inactivating proteins of plant origin, such as ricin, abrin, gelonin, and a number of others, and bacterial toxins such as pseudomonas exotoxin and diphtheria toxin.

Another useful class is the one including taxol, taxotere, and taxoids. Specific examples include paclitaxel (taxol), its analog docetaxel (taxotere), and derivatives thereof. The first two are clinical drugs used in treating a number of tumors while the taxoids act to induce cell death by inhibiting the depolymerization of tubulin. Such agents are readily linked to antibodies through disulfide bonds without disadvantageous effects on binding specificity.

In one instance, a truncated Pseudomonas exotoxin was fused to an anti-CD22 variable fragment and used successfully to treat patients with chemotherapy-resistant hairy-cell leukemia. (see: Kreitman et al., Efficacy of the anti-CD22 recombinant immunotoxin BL22 in chemotherapy-resistant hairy-cell leukemia, *N Engl J Med*, 345: 241-247 (2001)) Conversely, the cancer-linked peptides of the present invention offer the opportunity to prepare antibodies, recombinant or otherwise, against the appropriate antigens to target solid tumors, preferably those of malignancies of prostate tissue, using the same or similar cytotoxic conjugates. Thus, many of the previously used immunoconjugates have been formed using antibodies against general antigenic sites linked to cancers whereas the antibodies formed using the peptides disclosed herein are more specific and target the antibody-cytotoxic agent to a particular tissue or organ, thus further reducing toxicity and other undesirable side effects.

In addition, the immunoconjugates formed using the antibodies prepared against the cancer-linked antigens disclosed herein can be formed by any type of chemical coupling. Thus, the cytotoxic agent of choice, along with the immunoglobulin, can be coupled by any type of chemical linkage, covalent or

non-covalent, including electrostatic linkage, to form the immunoconjugates of the present invention.

When used as immunoconjugates, the antitumor agents of the present invention represent a class of pro-drugs that are relatively non-toxic when first administered to an animal (due mostly to the stability of the immunoconjugate), such as a human patient, but which are targeted by the conjugated immunoglobulin to a cancer cell where they then exhibit good toxicity. The tumor-related, associated, or linked, antigens, preferably those presented herein, serve as targets for the antibodies (monoclonal, recombinant, and the like) specific for said antigens. The end result is the release of active cytotoxic agent inside the cell after binding of the immunoglobulin portion of the immunoconjugate.

The cited references describe a number of useful procedures for the chemical linkage of cytotoxic agents to immunoglobulins and the disclosures of all such references cited herein are hereby incorporated by reference in their entirety. For other reviews see Ghetie et al., Immunotoxins in the therapy of cancer: from bench to clinic, *Pharmacol Ther*, **63**: 209-234 (1994), Pietersz et al. The use of monoclonal antibody immunoconjugates in cancer therapy, *Adv Exp Med Biol*, **353**:169-179 (1994), and Pietersz, G. A. The linkage of cytotoxic drugs to monoclonal antibodies for the treatment of cancer, *Bioconjug Chem*, **1**:89-95 (1990).

Thus, the present invention provides highly useful cancer-associated antigens for generation of antibodies for linkage to a number of different cytotoxic agents which are already known to have some *in vitro* toxicity and possess chemical groups available for linkage to antibodies.

The present invention also relates to a process that comprises a method for producing a product comprising identifying an agent according to one of the disclosed processes for identifying such an agent (i.e., the therapeutic agents



identified according to the assay procedures disclosed herein) wherein said product is the data collected with respect to said agent as a result of said identification process, or assay, and wherein said data is sufficient to convey the chemical character and/or structure and/or properties of said agent. For example, the present invention specifically contemplates a situation whereby a user of an assay of the invention may use the assay to screen for compounds having the desired enzyme modulating activity and, having identified the compound, then conveys that information (i.e., information as to structure, dosage, etc) to another user who then utilizes the information to reproduce the agent and administer it for therapeutic or research purposes according to the invention. For example, the user of the assay (user 1) may screen a number of test compounds without knowing the structure or identity of the compounds (such as where a number of code numbers are used the first user is simply given samples labeled with said code numbers) and, after performing the screening process, using one or more assay processes of the present invention, then imparts to a second user (user 2), verbally or in writing or some equivalent fashion, sufficient information to identify the compounds having a particular modulating activity (for example, the code number with the corresponding results). This transmission of information from user 1 to user 2 is specifically contemplated by the present invention.

It should be cautioned that, in carrying out the procedures of the present invention as disclosed herein, whether to form immunoconjugates or screen for other antitumor agents using the genes and polypeptides disclosed herein, any reference to particular buffers, media, reagents, cells, culture conditions and the like are not intended to be limiting, but are to be read so as to include all related materials that one of ordinary skill in the art would recognize as being of interest or value in the particular context in which that discussion is presented. For example, it is often possible to substitute one buffer system or culture medium for another and still achieve similar, if not identical, results. Those of skill in the art will have sufficient knowledge of such systems and methodologies so as to be

able, without undue experimentation, to make such substitutions as will optimally serve their purposes in using the methods and procedures disclosed herein.

5 The present invention will now be further described by way of the following non-limiting example. In applying the disclosure of the example, it should be kept clearly in mind that other and different embodiments of the methods disclosed according to the present invention will no doubt suggest themselves to those of skill in the relevant art. The following example shows how a potential anti-neoplastic agent may be identified using one or more of the genes disclosed  
10 herein.

## EXAMPLE

### 15 Determination of Gene Inhibitory Activity of an Anti-neoplastic Agent

SW480 cells are grown to a density of  $10^5$  cells/cm<sup>2</sup> in Leibovitz's L-15 medium supplemented with 2 mM L-glutamine (90%) and 10% fetal bovine serum. The cells are collected after treatment with 0.25% trypsin, 0.02% EDTA at  
20 37°C for 2 to 5 minutes. The trypsinized cells are then diluted with 30 ml growth medium and plated at a density of 50,000 cells per well in a 96 well plate (100  $\mu$ l/well). The following day, cells are treated with either compound buffer alone, or compound buffer containing a chemical agent to be tested, for 24 hours. The media is then removed, the cells lysed and the RNA recovered using the  
25 RNAeasy reagents and protocol obtained from Qiagen. RNA is quantitated and 10 ng of sample in 1  $\mu$ l are added to 24  $\mu$ l of Taqman reaction mix containing 1X PCR buffer, RNAsin, reverse transcriptase, nucleoside triphosphates, amplitaq gold, Tween 20, glycerol, bovine serum albumin (BSA) and specific PCR primers and probes for a reference gene (18S RNA) and a test gene (Gene X). Reverse  
30 transcription is then carried out at 48°C for 30 minutes. The sample is then applied to a Perkin Elmer 7700 sequence detector and heat denatured for 10

minutes at 95°C. Amplification is performed through 40 cycles using 15 seconds annealing at 60°C followed by a 60 second extension at 72°C and 30 second denaturation at 95°C. Data files are then captured and the data analyzed with the appropriate baseline windows and thresholds.

5

The quantitative difference between the target and reference gene is then calculated and a relative expression value determined for all of the samples used. In this way, the ability of a chemotherapeutic agent to effectively and selectively reduce the activity of a cancer-specific gene is readily ascertained.

- 10 The overall expression of the cancer-specific gene, as modulated by one chemical agent relative to another, is also determined. Chemical agents having the most effect in reducing gene activity are thereby identified as the most anti-neoplastic.

15

## References:

- Walter A. Blättler and Ravi Chari: Drugs to enhance the therapeutic potency of anti-cancer antibodies: antibody-drug conjugates as tumor-activated prodrugs. In  
5 *Anticancer Agents - Frontiers in Cancer Chemotherapy* (Iwao Ojima, Gregory D. Vite, Karl-Heinz Altmann, Eds.), American Chemical Society, pp. 317-338 (2001).
- Dan L. Longo, Patricia L. Duffey, John G. Gribben, Elaine S. Jaffe, Brendan D. Curti, Barry L. Gause, John E. Janik, Virginia M. Braman, Dixie Esseltine,  
10 Wyndham H. Wilson, Dwight Kaufman, Robert E. Wittes, Lee M. Nadler, and Walter J. Urba: Combination chemotherapy followed by an Immunotoxin (Anti-B4-blocked Ricin) in patients with indolent lymphoma: results of a Phase II study. *Cancer J.* 6, 146-150 (2000).
- Walter A. Blättler and John M. Lambert: Preclinical immunotoxin development. In  
15 *Monoclonal Antibody-Based Therapy of Cancer* (M. Grossbard, Ed.), Marcel Dekker, Inc. NY, NY, pp. 1-22 (1998).
- Ravi V. J. Chari: Targeted delivery of chemotherapeutics: tumor-activated prodrug therapy. In *Advanced Drug Delivery Reviews*, Elsevier Science B.V., pp. 89-104 (1998).
- David T. Scadden, David P. Schenkein, Zale Bernstein, Barry Luskey, John  
20 Doweiko, Anil Tulpule, and Alexandra M. Levine: Immunotoxin combined with chemotherapy for patients with AIDS-related Non-Hodgkin's Lymphoma. *Cancer* 83, 2580-2587 (1998).
- Changnian Liu and Ravi VJ Chari: The development of antibody delivery systems to target cancer with highly potent maytansinoids. *Exp. Opin. Invest. Drugs* 6, 169-  
25 172 (1997).
- A. C. Goulet, Viktor S. Goldmacher, John M. Lambert, C. Baron, Dennis C. Roy and E. Kouassi: Conjugation of blocked ricin to an anti-CD19 monoclonal antibody increases antibody-induced cell calcium mobilization and CD19 internalization. *Blood* 90, 2364-2375 (1997).
- Changnian Liu, John M. Lambert, Beverly A. Teicher, Walter A. Blättler, and  
30 Rosemary O'Connor: Cure of multidrug-resistant human B-cell lymphoma xenografts by combinations of anti-B4-blocked ricin and chemotherapeutic drugs. *Blood* 87, 3892-3898 (1996).
- Rajeeva Singh, Lana Kats, Walter A. Blättler, and John M. Lambert: Formation of  
35 N-Substituted 2-Iminoethanolanes when amino groups in proteins and peptides are modified by 2-Iminoethanolane. *Anal. Biochem.* 236, 114-125 (1996).
- Changnian Liu, B. Mitra Tadayoni, Elizabeth A. Bourret, Kristin M. Mattocks, Susan M. Derr, Wayne C. Widdison, Nancy L. Kedersha, Pamela D. Ariniello, Victor S. Goldmacher, John M. Lambert, Walter A. Blättler, and Ravi V.J. Chari:

Eradication of large colon tumor xenografts by targeted delivery of maytansinoids. *Proc. Natl. Acad. Sci. USA* 93, 8618-8623 (1996).

Denis C. Roy, Sophie Ouellet, Christiane Le Houiller, Pamela D. Ariniello, Claude Perreault and John M. Lambert: Elimination of neuroblastoma and small-cell lung cancer cells with an anti-neural cell adhesion molecule immunotoxin. *J. Natl. Cancer Inst.* 88, 1136-1145 (1996).

Walter A. Blättler, Ravi V.J. Chari and John M. Lambert: Immunoconjugates. In *Cancer Therapeutics: Experimental and Clinical Agents*. (B. Teicher, Ed.), Humana Press, Totowa, NJ, pp. 371-394 (1996).

10 Michael L. Grossbard, John M. Lambert, Victor S. Goldmacher, Arnold S. Freedman, Jeanne Kinsella, Danny P. Duccello, Susan N. Rabinowe, Laura Elisea, Felice Carol, James A. Taylor, Walter A. Blättler, Carol L. Epstein, and Lee M. Nadler: Anti-B4-blocked Ricin: A phase I trial of 7 day continuous infusion in patients with B-cell neoplasms. *J. Clin. Oncol.* 11, 726-737 (1993).

15 Michael L. Grossbard, John G. Gribben, Arnold S. Freedman, John M. Lambert, Jeanne Kinsella, Susan N. Rabinowe, Laura Eliseo, James A. Taylor, Walter A. Blättler, Carol L. Epstein, and Lee M. Nadler: Adjuvant immunotoxin therapy with anti-B4-blocked ricin following autologous bone marrow transplantation for patients with B-cell Non-Hodgkin's lymphoma. *Blood* 81, 2263-2271 (1993).

20 Sudhir A. Shah, Patricia M. Halloran, Cynthia A. Ferris, Beth A. Levine, Elizabeth A. Bourret, Victor S. Goldmacher, and Walter A. Blättler: Anti-B4-blocked Ricin immunotoxin shows therapeutic efficacy in four different SCID mouse tumor models. *Cancer Res.* 53, 1360-1367 (1993).

25 Ravi V.J. Chari, Bridget A. Martell, Jonathan L. Gross, Sherilyn B. Cook, Sudhir A. Shah, Walter A. Blättler, Sara J. McKenzie, and Victor S. Goldmacher: Immunoconjugates containing novel maytansinoids: promising anti-cancer drugs. *Cancer Res.* 52, 127-131 (1992).

30 John M. Lambert, Peter D. Senter, Annie Yau-Young, Walter A. Blättler, and Victor S. Goldmacher: Purified immunotoxins that are reactive with human lymphoid cells. *J. Biol. Chem.* 250, 12035-12041 (1985).

WHAT IS CLAIMED IS:

1. A process for identifying an agent that modulates the activity of a cancer-related gene comprising:

5 (a) contacting a compound with a cell containing a gene that corresponds to the polynucleotide having the sequence of SEQ ID NO: 2 or 4 and under conditions promoting the expression of said gene; and

(b) detecting a difference in expression of said gene relative to when said compound is not present

10 thereby identifying an agent that modulates the activity of a cancer-related gene.

2 The process of claim 1 wherein said gene has the sequence of SEQ ID NO: 2 or 4.

15

3. The process of claim 1 or 2 wherein the cell is a cancer cell and the difference in expression is a decrease in expression.

4. The process of claim 3 wherein said cancer cell is a prostate cancer  
20 cell.

5. A process for identifying an anti-neoplastic agent comprising contacting a cell exhibiting neoplastic activity with a compound first identified as a cancer related gene modulator using a process of one of claims 1, 2, 3 or 4 and  
25 determining a decrease in said neoplastic activity after said contacting compared to when said contacting does not occur.

6. The process of claim 5 wherein said neoplastic activity is accelerated cellular replication.

30

7. The process of claim 5 wherein said decrease in neoplastic activity results from the death of the cell.

8. A process for identifying an anti-neoplastic agent comprising  
5 administering to an animal exhibiting a cancer condition an effective amount of an agent first identified according to a process of one of claims 1, 2, 3, 4, 5, 6 or 7 and detecting a decrease in said cancerous condition.

9. A process for determining the cancerous status of a cell, comprising  
10 determining an increase in the level of expression in said cell of a gene that corresponds to a polynucleotide having the sequence of SEQ ID NO: 2 or 4 wherein an elevated expression relative to a known non-cancerous cell indicates a cancerous state or potentially cancerous state.

15 10. The process of claim 9 wherein said elevated expression is due to an increased copy number.

11. An isolated polypeptide comprising an amino acid sequence homologous to the amino acid sequence of SEQ ID NO: 3 or 5 wherein any  
20 difference between said amino acid sequence and the sequence of SEQ ID NO: 3 or 5 is due solely to conservative amino acid substitutions and wherein said isolated polypeptide comprises at least one immunogenic fragment.

12. An isolated polypeptide comprising the amino acid sequence of SEQ  
25 ID NO: 3 or 5.

13. An antibody that reacts with a polypeptide comprising the amino acid sequence of SEQ ID NO: 3 or 5.

14. The antibody of claim 13 wherein said antibody is a recombinant  
30 antibody.

15. The antibody of claim 13 wherein said antibody is a synthetic antibody.
16. The antibody of claim 13 wherein said antibody is a humanized antibody.
- 5 17. An immunoconjugate comprising the antibody of claim 13 and a cytotoxic agent.
- 10 18. The antibody of claim 17 wherein said cytotoxic agent is a member selected from the group consisting of a calicheamicin, a maytansinoid, an adozelesin, a cytotoxic protein, a taxol, a taxotere, a taxoid and DC1.
- 15 19. The immunoconjugate of claim 18 wherein said calicheamicin is calicheamicin  $\gamma_1^I$ , N-acetyl gamma calicheamicin dimethyl hydrazide or calicheamicin  $\theta_1^I$ .
20. The immunoconjugate of claim 18 wherein said maytansinoid is DM1.
21. The immunoconjugate of claim 18 wherein said cytotoxic protein is ricin, abrin, gelonin, pseudomonas exotoxin or diphtheria toxin.
22. The immunoconjugate of claim 18 wherein said taxol is paclitaxel.
23. The immunoconjugate of claim 18 wherein said taxotere is docetaxel.
- 25 24. A process for treating cancer comprising contacting a cancerous cell *in vivo* with an agent having activity against an expression product encoded by the gene sequence of SEQ ID NO: 2 or 4.
- 30 25. The process of claim 24 wherein said agent is an antibody of claim 13
- 16.



26. The process of claim 24 wherein said agent is an immunoconjugate of claim 17.

5           27. An immunogenic composition comprising a polypeptide of claim 11.

28. An immunogenic composition comprising a polypeptide of claim 12.

10           29. The process of claim 24 wherein said cancer is prostate cancer.

30. A process for treating cancer in an animal afflicted therewith comprising administering to said animal an amount of an immunogenic composition of claim 27 sufficient to elicit the production of cytotoxic T lymphocytes specific for the polypeptide of claim 11.

15           31. A process for treating cancer in an animal afflicted therewith comprising administering to said animal an amount of an immunogenic composition of claim 28 sufficient to elicit the production of cytotoxic T lymphocytes specific for the polypeptide of claim 12.

20           32. A process for treating a cancerous condition in an animal afflicted therewith comprising administering to said animal a therapeutically effective amount of an agent first identified as having anti-neoplastic activity using the process of claim 8.

25           33. A process for protecting an animal against cancer comprising administering to an animal at risk of developing cancer a therapeutically effective amount of an agent first identified as having anti-neoplastic activity using the process of claim 8.

30           34. The process of claim 30, 31, 32 or 33 wherein said animal is a human being.

35. The process of claim 30, 31, 32 or 33 wherein said cancer is prostate cancer.

- 5           36. A method for producing a product comprising identifying an agent according to the process of claim 1, 2, 3, 4, 5, 6, 7 or 8 wherein said product is the data collected with respect to said agent as a result of said process and wherein said data is sufficient to convey the chemical structure and/or properties of said agent.

## ABSTRACT

Cancer-linked gene sequences, and derived amino acid sequences, are disclosed along with processes for assaying potential antitumor agents based on  
5 their modulation of the expression of these cancer-linked genes. Also disclosed are antibodies that react with the disclosed polypeptides and methods of using the antibodies to treat cancerous conditions, such as by using the antibody to target cancerous cells *in vivo* for purposes of delivering therapeutic agents thereto. Also described are methods of diagnosing using the gene sequences.

10